2019

MARYLAND ANNUAL HIV EPIDEMIOLOGICAL PROFILE

Data reported through June 30, 2020

CENTER FOR HIV SURVEILLANCE, EPIDEMIOLOGY AND EVALUATION

Infectious Disease Prevention and Health Services Bureau Prevention and Health Promotion Administration



Acknowledgements

This report is published by The Center for HIV Surveillance, Epidemiology and Evaluation (CHSEE). This report would not have been possible without the cooperation, dedication and hard work of HIV health care providers, local health departments, community groups, researchers, and members of the community.

This report was prepared by the following CHSEE staff: Alyssa Arey, Morgan Boyer, James Carrier, Colin Flynn, Molly Gribbin, Trevor Hsu, Amber Lee, Michelle Ryan, and Jami Stockdale.

Maryland Department of Health Non-Discrimination Statement

The Maryland Department of Health (MDH) complies with applicable Federal civil right laws and does not discriminate on the basis of race, color, national origin, age, disability in its health programs and activities.

English

Help is available in your language: 410-767-5227 (TTY: 800-735-2258). These services are available for free.

Español/Spanish

Hay ayuda disponible en su idioma: 410-767-5227 (TTY: 800-735-2258). Estos servicios están disponibles gratis.

中文/Chinese

用您的语言为您提供帮助: 410-767-5227 (TTY: 800-735-2258). 这些服务都是免费的

Suggested Citation

Maryland Annual HIV Epidemiological Profile 2019. Center for HIV Surveillance, Epidemiology and Evaluation, Maryland Department of Health, Baltimore, MD. 2020.

On the Web

https://phpa.health.maryland.gov/OIDEOR/CHSE/SiteAssets/Pages/statistics/Maryland-Annual-HIV-Epidemiological-Profile-2019

Additional Information and Reporting Instructions

https://phpa.health.maryland.gov/OIDEOR/CHSE/pages/Home.aspx

Email: mdh.hivdatarequest@maryland.gov

Table of Contents

Section I – Summary of Analyses and Technical Notes	6
Section II – Trends 1985-2019	10
Figure 1 – Trends in Reported HIV and AIDS Diagnoses and Deaths, Among Residents at Diagnosis, 1985-2019,	
Reported through June 30, 2020	10
Figure 1A – Trends in Rates of Reported HIV and AIDS Diagnoses, Among Residents at Diagnosis Aged 13+, 2010-	
2019, Reported through June 30, 2020	11
Figure 2 – Trends in People Living with Diagnosed HIV, Among Residents at Diagnosis, 1985-2019, Reported	
through June 30, 2020	12
Figure 3 – Trends in People Aged 13+ Living with Diagnosed HIV, Among Current Residents, 2010-2019, Reported	
through June 30, 2020	13
Figure 4 – Trends in Reported HIV Diagnoses, Among Residents at Diagnosis Aged 13+, by Sex at Birth, 1985-	
2019, Reported through June 30, 2020	14
Figure 4A – Trends in Rates of Reported HIV Diagnoses, Among Residents at Diagnosis Aged 13+, by Sex at Birth,	
2010-2019, Reported through June 30, 2020	16
Figure 5 – Trends in Reported HIV Diagnoses, Among Residents at Diagnosis Aged 13+, by Race/Ethnicity, 1985-	
2019, Reported through June 30, 2020	17
Figure 5A – Trends in Rates of Reported HIV Diagnoses, Among Residents at Diagnosis Aged 13+, by	
Race/Ethnicity, 2010-2019, Reported through June 30, 2020	19
Figure 6 – Trends in Reported HIV Diagnoses, Among Residents at Diagnosis Aged 13+, 1985-2019, Reported	
through June 30, 2020	20
Figure 6A – Trends in Rates of Reported HIV Diagnoses, Among Residents at Diagnosis Aged 13+, by Age at	
Diagnosis, 2010-2019, Reported through June 30, 2020	22
Figure 7 – Trends in Reported HIV Diagnoses, Among Residents at Diagnosis Aged 13+, by Estimated Exposure	
Category, 1985-2019, Reported through June 30, 2020	23
Section III – Continuum of Care Cascades	25
Figure 8 – Prevalence-Based Estimated 2019 HIV Continuum of Care Cascades, Among Current Maryland	
Residents Aged 13+, Reported through June 30, 2020	25
Figure 8A – Prevalence-Based Estimated 2019 HIV Continuum of Care Cascades, Among Current Maryland	
Residents Aged 13+, by Sex at Birth, Reported through June 30, 2020	26
Figure 8B – Prevalence-Based Estimated 2019 HIV Continuum of Care Cascades, Among Current Maryland	
Residents Aged 13+, by Race/Ethnicity, Reported through June 30, 2020	27
Figure 8C – Prevalence-Based Estimated 2019 HIV Continuum of Care Cascades, Among Current Maryland	
Residents Aged 13+, by Age on December 31, 2019, Reported through June 30, 2020	28
Figure 8D – Prevalence-Based Estimated 2019 HIV Continuum of Care Cascades, Among Current Maryland	
Residents Aged 13+, by Estimated Exposure Category, Reported through June 30, 2020	29
Figure 9 – Diagnosis-Based Estimated 2019 HIV Continuum of Care Cascades, Among Current Maryland	
Residents Aged 13+, Reported through June 30, 2020	31
Section IV – By Jurisdiction	32
Table 1 – Reported HIV Diagnoses and Reported AIDS Diagnoses, Among People Aged 13+ During 2019, by	
Jurisdiction of Residence at Diagnosis, Reported through June 30, 2020	33
Table 2 – Reported HIV Diagnoses, Among People Aged 13+ During 2019, Linked to Care, Late Diagnosis, First	
CD4 Test Result, and Mean Years from HIV Diagnosis by Jurisdiction of Residence at Diagnosis,	
Reported through June 30, 2020	34

Table 3 – People Aged 13+ Living with Diagnosed HIV, Alive on December 31, 2019, by Jurisdiction of Residence	
at Diagnosis and Current Residence, Reported through June 30, 2020	35
Table 4 – CD4 Test Results During 2019 for People Aged 13+ Living with Diagnosed HIV, Alive on December 31,	
2019, by Jurisdiction of Current Residence, Reported through June 30, 2020	36
Table 5 – Viral Load Test Results During 2019 for People Aged 13+ Living with Diagnosed HIV, Alive on December	r
31, 2019, by Jurisdiction of Current Residence, Reported through June 30, 2020	37
Section V – By Age	38
Table 6 – Reported HIV Diagnoses During 2019, Linked to Care, Late Diagnosis, and First CD4 Test Result by Age	
at HIV Diagnosis, Reported through June 30, 2020	
Table 7 – People Living with Diagnosed HIV, Alive on December 31, 2019, by Age on December 31, 2019,	
Reported through June 30, 2020	38
Table 8 – CD4 Test Results During 2019 for People Aged 13+ Living with Diagnosed HIV, Alive on December 31,	
2019, by Age on December 31, 2019, Reported through June 30, 2020	38
Table 9 – Viral Load Test Results During 2019 for People Living with Diagnosed HIV, Alive on December 31, 2019	
by Age on December 31, 2019, Reported through June 30, 2020	-
Section VI – By Demographics	40
Table 10 – Reported HIV Diagnoses, Among People Aged 13+ During 2019, Linked to Care, Late Diagnosis, and	
First CD4 Test Result by Sex at Birth, Gender, Race/Ethnicity, and Country of Birth, Reported	40
through June 30, 2020	40
Table 11 – Reported HIV Diagnoses, Among People Aged 13+ During 2019, by Sex at Birth and Race/Ethnicity,	11
Reported through June 30, 2020	41
Table 12 – People Aged 13+ Living with Diagnosed HIV Alive on December 31, 2019, by Sex at Birth, Gender,	42
Race/Ethnicity, and Country of Birth, Reported through June 30, 2020	42
Table 13 – People Aged 13+ Living with Diagnosed HIV Alive on December 31, 2019, by Sex at Birth and	42
Race/Ethnicity, Reported through June 30, 2020	
Table 14 – CD4 Test Results During 2019 for People Aged 13+ Living with Diagnosed HIV, Alive on December 31,	
2019, by Sex at Birth, Gender, Race/Ethnicity and Country of Birth, Reported through June 30, 202	J44
Table 15 – Viral Load Test Results During 2019 for People Aged 13+ Living with Diagnosed HIV, Alive on	
December 31, 2019, by Sex at Birth, Gender, Race/Ethnicity and Country of Birth, Reported throug	
June 30, 2020	45
Table 16 – Males Living with Diagnosed HIV by Age on December 31, 2019 and Race/Ethnicity, Alive on	4.0
December 31, 2019, and Reported through June 30, 2020	46
Table 17 – Females Living with Diagnosed HIV by Age on December 31, 2019 and Race/Ethnicity, Alive on	47
December 31, 2019, and Reported through June 30, 2020	
Section VII – By HIV Exposure Category	48
Table 18 – Reported HIV Diagnoses, Among People Aged 13+ During 2019, Linked to Care, Late Diagnosis, and	
First CD4 Test Result, by Estimated or Reported Exposure Category and Sex at Birth, Reported	
through June 30, 2020	48
Table 19 – People Aged 13+ Living with Diagnosed HIV, Alive on December 31, 2019, by Estimated or Reported	
Exposure Category and Sex at Birth, Reported through June 30, 2020	49
Table 20 – People Aged 13+ Living with Diagnosed HIV by Estimated or Reported Exposure Category,	
Race/Ethnicity, and Sex at Birth, Alive on December 31, 2019, and Reported through June 30, 2020	
Table 21 – CD4 Test Results During 2019 for People Aged 13+ Living with Diagnosed HIV, Alive on December 31,	
2019. by Estimated or Reported Exposure Category, Reported through June 30, 2020	51

Table 22 – Viral Load Test Results During 2019 for People Aged 13+ Living with Diagnosed HIV, Alive on	
December 31, 2019, by Estimated or Reported Exposure Category, Reported through June 30, 2020	51
Section VIII – Disease Progression and Mortality	52
Figure 10 – Trends in Disease Progression by Year of HIV Diagnosis, 1985-2018, Reported through June 30, 2020	52
Figure 11 – Trends in Survival Times by Year of AIDS Diagnosis, 1985-2018, Reported through June 30, 2020	54
Section IX – Pediatrics	56
Figure 12 – Trends in Reported HIV Diagnoses Among People Aged <13, 1985-2019, Reported through June 30,	
2020	56
Table 23 – People Aged <13 Living with Diagnosed HIV, Alive on December 31, 2019, by Jurisdiction of Current	
Residence, Reported through June 30, 2020	57
Table 24 – People Aged <13 Living with Diagnosed HIV, Alive on December 31, 2019, by Sex at Birth and	
Race/Ethnicity, Reported through June 30, 2020	58
Table 25 – Perinatal HIV Transmissions by Year of Birth, 2010-2019	58
Glossary of Terms	59

Section I – Summary of Analyses and Technical Notes

Highlights of Analyses

The Center for HIV Surveillance, Epidemiology and Evaluation in the Maryland Department of Health is responsible for the collection, analysis, and dissemination of HIV surveillance data for the state of Maryland. The annual epidemiological profiles, fact sheets, maps and other reports attempt to summarize information about the HIV epidemic in Maryland.

All rates are per 100,000 population.

HIV Diagnoses

During 2019, there were 931 reported HIV diagnoses in Maryland. Data presented below describe HIV diagnoses in Maryland, among people age 13 years and older, during 2019.

- <u>Geographic distribution:</u> Prince George's County accounted for the highest number (289, 31.0%) of HIV diagnoses. The rate was highest in Baltimore City (39.7).
- Age: Persons aged 20-39 years accounted for the highest number (566, 60.8%) of HIV diagnoses. The highest rate was among 20-29-year-olds (41.5).
- Sex at Birth: Males accounted for the highest number (662, 71.1%) of HIV diagnoses. The rate among males was 27.1 and 10.2 among females.
- Race/Ethnicity: Non-Hispanic Blacks accounted for the highest number (690, 74.1%) and the highest rate (45.4) of HIV diagnoses.
- Exposure Category: Male-to-male sexual contact accounted for the highest number (477, 51.2%) of diagnoses.

People living with Diagnosed HIV

At the end of 2019, there were 31,630 people living with diagnosed HIV in Maryland. Data presented below describe people living with diagnosed HIV, aged 13 years and older, in Maryland.

- <u>Geographic distribution:</u> Baltimore City accounted for the highest number (10,600, 33.5%) of people living with diagnosed HIV, as well as the highest rate (2,102.1).
- Age: Persons aged 50+ years accounted for the highest number (16,181, 51.1%) of people living with diagnosed HIV. The highest rate was among 50-59-year-olds (1,122.9).
- <u>Sex at Birth:</u> Males accounted for the highest number (20,893, 66.1%) of people living with diagnosed HIV. The rate among males was 856.4 and 405.6 among females.
- Race/Ethnicity: Non-Hispanic Blacks accounted for the highest number (23,578, 74.5%) of the people living with diagnosed HIV, as well as the highest rate (1,552.1).
- Exposure Category: Male-to-male sexual contact accounted for the highest number (12,576, 39.8%) of the total people living with diagnosed HIV.

Engagement in HIV Care

- <u>HIV diagnosed:</u> Among people living with HIV in Maryland, it is estimated that 89.2% have been diagnosed, while an estimated 3,830 people with HIV in Maryland remain undiagnosed.
- <u>Linked to care:</u> During 2019, 86.9% of reported HIV diagnoses were linked to HIV care within one month of their HIV diagnosis.
- Retained in care: Among people living with diagnosed HIV, 80.5% were retained in care during 2019.
- Viral suppression: In 2019, 69.8% of people living with diagnosed HIV achieved viral suppression.

Technical Notes

HIV/AIDS Reporting

The Maryland HIV/AIDS Reporting Act of 2007 requires that people living with HIV be reported by name to the Maryland Department of Health. As per the reporting requirements of Health-General Articles 18-201.1, 18-202.1, and 18-205 of the Annotated Code of Maryland, as specified in the Code of Maryland Regulations (COMAR) 10.18.02, physicians are required to report patients in their care with diagnoses of HIV or AIDS and infants born to mothers living with HIV, clinical and infection control practitioners in medical and correctional facilities are required to report patients in the care of the institution with diagnoses of HIV or AIDS, and laboratory directors are required to report all HIV diagnostic and HIV-related monitoring test results, including HIV negative test results.

Report Changes

In order to be responsive to changing data needs, the following updates have been made to this report:

- The Background Information section has been replaced with the Summary of Analyses and Technical Notes section.
- Tables describing new HIV diagnoses by geography have been restructured. All new HIV and AIDS diagnoses will appear in Table 1. All laboratory data and outcomes for new HIV diagnoses will be described in Table 2. In the county level reports, only ZIP codes with at least five new HIV diagnoses during the specified year will be included in tables describing laboratory data and outcomes.
- Tables describing people living with diagnosed HIV by sex at birth and race/ethnicity have been added to the demographics section (Section VI).
- Mean years from HIV to AIDS diagnosis and the percentage of late HIV diagnoses among AIDS diagnoses is no longer presented in tables. Aggregate disease progression data by year is presented in Figure 10.

- The population pyramids figure describing people living with diagnosed HIV by race/ethnicity, sex at birth, and age is no longer included in this report.
- The treemap figure presenting data by sex at birth, race/ethnicity, and exposure category is no longer included in this report.

Data Specifications

Surveillance is the ongoing systematic collection, analysis, interpretation, and dissemination of data. Data are only available for people that have been diagnosed with HIV by a health care provider, were residents of Maryland at the time of diagnosis or are current residents of Maryland as of December 31, 2019, are receiving medical care (often only at facilities in Maryland), and only includes information that has been reported to the health department.

Data presented were reported to the Maryland Department of Health through June 30, 2020. Reported new diagnoses and estimates of people living with diagnosed HIV are as of December 31, 2019.

This report presents data with a six-month reporting delay. This lag allows for time to report case and laboratory data and to complete investigations. Due to reporting delays, estimates by exposure category and mortality are preliminary.

Please read all table titles and footnotes carefully to ensure a complete understanding of the displayed data.

New HIV Infections

The Centers for Disease Control and Prevention (CDC), using national surveillance data, estimates that the number of people living with undiagnosed HIV infection, among people living with HIV, is 13.8 percent in the United States. Applying the same estimation method to Maryland surveillance data, the estimated number of people living with undiagnosed HIV infection in Maryland is 10.8 percent in 2018.

¹Centers for Disease Control and Prevention. Estimated HIV incidence and prevalence in the United States, 2014–2018. HIV Surveillance Supplemental Report 2020;25(No. 1). http://www.cdc.gov/hiv/library/reports/hiv-surveillance.html. Published May 2020.

AIDS Diagnoses

CD4+ T-lymphocyte tests are measures of a person's immune system function. An adult living with diagnosed HIV is considered to have AIDS if they have less than 200 CD4+ cells per microliter of blood or if the percent of T-Lymphocyte cells that are CD4+ is less than 14 percent.

Mortality

People are assumed to be alive unless reported as deceased to the Maryland Department of Health. Caution should be exercised when interpreting mortality estimates for recent years due to delays in reporting.

Engagement in HIV Care

Treatment recommendations are that a person in HIV medical care should have their CD4 and viral load levels measured regularly, at least once per year. The presence of these lab tests is used as an indicator that someone has been "linked to care" after diagnosis or is "retained in care."

Viral Suppression

Viral load tests are measures of the amount of HIV in a person's body. The goal of HIV treatment is to have a low number of copies of virus per milliliter of blood, below what the test can measure, which is called an undetectable level. Low levels of viral load, such as less than 200 copies per milliliter of blood, is known as viral suppression. Estimates of viral suppression among people living with diagnosed HIV are inclusive of people diagnosed in the specified year; therefore, they may not have had sufficient time to engage in HIV care and/or achieve viral suppression.

Geographic Distribution

Residence at diagnosis is used exclusively to describe new HIV and AIDS diagnoses. Estimates describing the number of people living with diagnosed HIV use current residence. Current residence estimates are restricted to people for which there is a case report form or laboratory test reported since January 1, 2009. Restricting address data to recent years presents the most accurate data available and helps to account for people that may have moved out of state whose data would no longer be reported in Maryland. However, current residence data excludes people that may still be residents of Maryland

but have not received any HIV care since prior to 2009. In addition, residence is dynamic and people living with diagnosed HIV may have resided at multiple addresses that cannot all be represented in single time point estimates.

Residents of Correctional Facilities

People living with diagnosed HIV who were reported to be residing in a state or federal prison will be described for Maryland only. For surveillance purposes, pre- or post-incarceration residential addresses are used for people living with diagnosed HIV who were incarcerated in local jails and detention centers.

Gender Identity

This epidemiological profile describes people by gender identity. Gender identity is one's internal understanding of one's own gender or the gender with which a person identifies. Historically, it has been difficult to present data by gender identity. No reliable system exists for collecting or reporting gender identity and some agencies do not collect or have complete data on gender identity. This report is likely underreporting the number gender diverse people (defined as people who have a gender identity or gender expression that differs from their assigned sex at birth) living with diagnosed HIV because of challenges in accurately identifying and reporting gender identity in HIV surveillance.

Race/Ethnicity

The completeness of reporting for race/ethnicity is variable in surveillance data. Definitions of race/ethnicity categories may vary across agencies and some agencies do not collect or have complete data on race/ethnicity.

Country of Birth

The completeness of reporting a person's country of birth is variable in surveillance data. Not all agencies collect or have complete data on nativity. In addition, some individuals may immigrate to the U.S. with HIV. HIV testing data from other countries is often unavailable for these people; therefore, they are counted as a new diagnosis following their first HIV related lab test in the United States, even if they were first diagnosed elsewhere.

HIV Exposure Categories

The CDC developed a hierarchy of exposure categories for surveillance purposes. Persons with more than one reported risk factor are classified in the exposure category listed first in the hierarchy and therefore counted only once. Men who report a history of sexual contact with other men and report injection drug use constitutes a separate transmission category.

Sources of Data

Information on HIV and AIDS diagnoses, including residence, age, race/ethnicity, sex at birth, current gender, country of birth, vital status, HIV exposure category, and CD4 and HIV viral load test results are from the Maryland Department of Health's Enhanced HIV/AIDS Reporting System (eHARS), as of June 30, 2020.

Population data by sex, age, and race/ethnicity are from the July 1, 2019 U.S. Census Estimates. Population data by country-of-birth are from the Census Bureau's 2018 American Community Survey. This produces a difference in total population due to the different source. Due to estimation limitations, some population totals may not equal the sum of its components. When needed, age groups were divided by assuming uniform age distribution within the age group. Non-Hispanic multiple race and Non-Hispanic some other race from the Census were combined into one group.

Tabulation of Column Totals

Numbers in figures, tables and generally in the text have been rounded. Discrepancies in tables between totals and sums of components are due to rounding.

Data Suppression

In order to protect the confidentiality of people living with diagnosed HIV, data are suppressed in the following instances:

- Data describing a demographic group or geographic area (e.g. ZIP code) with a population less than 1,000 people.
- All clinical/laboratory information if it is describing less than five people.
- If any cell is suppressed, additional cells are also suppressed as necessary to prevent back calculation of the suppressed cell(s).

Tables presenting clinical/laboratory information for reported new diagnoses by ZIP code will present only ZIP codes with at least five people.

Exposure/risk data is not suppressed due to statistical adjustments resulting in the presentation of estimated or probable risk.


Multiple Imputation

Not all exposure data is reported and is therefore missing in some instances. Multiple imputation is a statistical method by which missing exposure categories are replaced with estimated or probable exposure categories. Perinatal transmission and other exposure categories are never estimated or adjusted and depict the reported exposure category.

Section II – Trends 1985-2019

<u>Figure 1 – Trends in Reported HIV and AIDS Diagnoses and Deaths, Among Residents at Diagnosis, 1985-2019, Reported through June 30, 2020</u>

People Diagnosed with HIV (Reported HIV Diagnoses) by Year of HIV Diagnosis, People Diagnosed with HIV with an AIDS Diagnosis (Reported AIDS Diagnoses) by Year of AIDS Diagnosis, and People Diagnosed with HIV with an AIDS Diagnosis Reported to have Died of Any Cause (Reported AIDS Deaths) by Year of Death, from 1985 through 2019, as Reported through June 30, 2020





Year of Diagnosis or Death	Reported HIV Diagnoses	Reported AIDS Diagnoses	Reported AIDS Deaths	Year of Diagnosis or Death	Reported HIV Diagnoses	Reported AIDS Diagnoses	Reported AIDS Deaths
<1985	386	149	72	2002	2,036	1,582	847
1985	724	221	121	2003	1,917	1,521	883
1986	947	320	180	2004	1,960	1,490	750
1987	1,259	510	279	2005	2,001	1,430	812
1988	1,707	712	360	2006	2,145	1,302	683
1989	2,165	956	544	2007	2,188	1,085	693
1990	2,285	1,197	662	2008	2,056	1,057	625
1991	2,605	1,504	846	2009	1,706	919	588
1992	2,459	1,990	1,001	2010	1,746	1,020	533
1993	2,218	2,291	1,227	2011	1,433	825	490
1994	2,038	2,206	1,474	2012	1,353	882	485
1995	2,008	2,182	1,627	2013	1,313	866	486
1996	1,867	1,960	1,307	2014	1,262	644	489
1997	1,888	1,695	802	2015	1,189	637	489
1998	1,847	1,545	751	2016	1,125	600	509
1999	1,810	1,543	785	2017	1,053	590	463
2000	1,721	1,423	720	2018*	1,024	518	491
2001	1,824	1,546	806	2019*	931	488	232
				Total	60,196	41,406	24,112

^{*2018-2019} reported AIDS deaths are preliminary

<u>Figure 1A – Trends in Rates of Reported HIV and AIDS Diagnoses, Among Residents at Diagnosis Aged 13+, 2010-2019, Reported through June 30, 2020</u>

Rates of Reported HIV and AIDS Diagnoses, Age 13+ at HIV Diagnosis, by Year of HIV Diagnosis from 2010 through 2019, as Reported through June 30, 2020






		Reported Diagnoses Among People Aged 13+						
Year of Diagnosis	Population Age 13+	ulation Age 13+ HIV Diagnoses			ses			
		Number	Rate	Number	Rate			
2010	4,829,275	1,743	36.1	1,011	20.9			
2011	4,878,213	1,430	29.3	817	16.7			
2012	4,923,893	1,350	27.4	877	17.8			
2013	4,959,623	1,310	26.4	855	17.2			
2014	4,990,288	1,256	25.2	639	12.8			
2015	5,019,656	1,189	23.7	628	12.5			
2016	5,038,152	1,121	22.3	593	11.8			
2017	5,059,735	1,050	20.8	585	11.6			
2018	5,074,057	1,021	20.1	509	10.0			
2019	5,086,800	931	18.3	480	9.4			

<u>Figure 2 – Trends in People Living with Diagnosed HIV, Among Residents at Diagnosis,</u> 1985-2019, Reported through June 30, 2020

People Living with Diagnosed HIV in Maryland and Not Reported to have Died as of December 31st of Each Year (People who were Diagnosed with HIV or AIDS in Maryland and/or People who are Currently Living with Diagnosed HIV in Maryland) from 1985 through 2019, as Reported through June 30, 2020




	People Living with	n Diagnosed HIV		People Living with	Diagnosed HIV
Year	Diagnosed with HIV or AIDS in Maryland	Current Residents	Year	Diagnosed with HIV or AIDS in Maryland	Current Residents
1985	917		2003	18,965	
1986	1,678		2004	20,089	
1987	2,660		2005	21,204	
1988	3,990		2006	22,564	
1989	5,579		2007	23,967	
1990	7,172		2008	25,282	
1991	8,868		2009	26,279	26,214
1992	10,238		2010	27,406	29,307
1993	11,105		2011	28,262	30,328
1994	11,568		2012	29,019	31,023
1995	11,847		2013	29,762	31,418
1996	12,299		2014	30,456	32,017
1997	13,299		2015	31,066	32,730
1998	14,286		2016	31,580	32,627
1999	15,175		2017	32,050	31,611
2000	16,072		2018	32,452	31,580
2001	16,984		2019	33,101	31,680
2002	18,073				

Note. Current residence for people living with diagnosed HIV is unavailable prior to 2009.

<u>Figure 3 – Trends in People Aged 13+ Living with Diagnosed HIV, Among Current Residents,</u> 2010-2019, Reported through June 30, 2020


People Living with Diagnosed HIV in Maryland and Not Reported to have Died as of December 31st of Each Year from 2010 through 2019, as Reported through June 30, 2020




		Maryland Residents Aged 13+ as of December 31st of the Specified Year							
Year	Population Age 13+	Diagnosed with HIV in Maryland Diagnosed with HIV in Outside of Maryland		Total	Rate				
2010	4,829,275	23,910	5,259	29,169	604.0				
2011	4,878,213	24,697	5,507	30,204	619.2				
2012	4,923,893	25,268	5,644	30,912	627.8				
2013	4,959,623	25,632	5,691	31,323	631.6				
2014	4,990,288	26,076	5,856	31,932	639.9				
2015	5,019,656	26,510	6,148	32,658	650.6				
2016	5,038,152	26,452	6,098	32,550	646.1				
2017	5,059,735	26,013	5,527	31,540	623.4				
2018	5,074,057	26,051	5,469	31,520	621.2				
2019	5,086,800	26,185	5,445	31,630	621.8				


<u>Figure 4 – Trends in Reported HIV Diagnoses, Among Residents at Diagnosis Aged 13+, by</u> Sex at Birth, 1985-2019, Reported through June 30, 2020

Number and Percent by Sex at Birth of People Diagnosed with HIV, Age 13+ at HIV Diagnosis, by Year of HIV Diagnosis from 1985 through 2019, as Reported through June 30, 2020





Reported HIV Diagnoses Among People Aged 13+ by Sex at Birth, 1985-2019, Reported through June 30, 2020




Reported HIV Diagnoses Among People Aged 13+

Year of HIV Diagnosis			Sex at Bi	rth		
real of filv Diagnosis	Number	Male		Female		
		Number	Percent	Number	Percent	
<1985	381	338	88.7%	43	11.3%	
1985	711	601	84.5%	110	15.5%	
1986	932	766	82.2%	166	17.8%	
1987	1,237	1,005	81.2%	232	18.8%	
1988	1,679	1,316	78.4%	363	21.6%	
1989	2,128	1,638	77.0%	490	23.0%	
1990	2,228	1,676	75.2%	552	24.8%	
1991	2,546	1,865	73.3%	681	26.7%	
1992	2,394	1,733	72.4%	661	27.6%	
1993	2,162	1,536	71.0%	626	29.0%	
1994	1,997	1,403	70.3%	594	29.7%	
1995	1,979	1,381	69.8%	598	30.2%	
1996	1,830	1,268	69.3%	562	30.7%	
1997	1,870	1,205	64.4%	665	35.6%	
1998	1,827	1,201	65.7%	626	34.3%	
1999	1,786	1,212	67.9%	574	32.1%	
2000	1,705	1,076	63.1%	629	36.9%	
2001	1,810	1,148	63.4%	662	36.6%	
2002	2,022	1,222	60.4%	800	39.6%	
2003	1,911	1,206	63.1%	705	36.9%	
2004	1,949	1,203	61.7%	746	38.3%	
2005	1,993	1,250	62.7%	743	37.3%	
2006	2,131	1,351	63.4%	780	36.6%	
2007	2,144	1,375	64.1%	769	35.9%	
2008	2,041	1,336	65.5%	705	34.5%	
2009	1,696	1,125	66.3%	571	33.7%	
2010	1,743	1,182	67.8%	561	32.2%	
2011	1,430	1,008	70.5%	422	29.5%	
2012	1,350	980	72.6%	370	27.4%	
2013	1,310	941	71.8%	369	28.2%	
2014	1,256	892	71.0%	364	29.0%	
2015	1,189	867	72.9%	322	27.1%	
2016	1,121	807	72.0%	314	28.0%	
2017	1,050	757	72.1%	293	27.9%	
2018	1,021	738	72.3%	283	27.7%	
2019	931	662	71.1%	269	28.9%	
Total	59,490	41,270	69.4%	18,220	30.6%	

<u>Figure 4A – Trends in Rates of Reported HIV Diagnoses, Among Residents at Diagnosis</u> <u>Aged 13+, by Sex at Birth, 2010-2019, Reported through June 30, 2020</u>


Rate by Sex at Birth of People Diagnosed with HIV, Age 13+ at HIV Diagnosis, by Year of HIV Diagnosis from 2010 through 2019, as Reported through June 30, 2020




Year of HIV Diagnosis	Number of HIV Diagnoses		Sex at Birt	th
rear of filv Diagnosis	Number of filv Diagnoses		Male	Female
		Population	2,311,027	2,518,248
2010	1,743	Reported Diagnoses	1,182	561
		Rate	51.1	22.3
		Population	2,336,414	2,541,799
2011	1,430	Reported Diagnoses	1,008	422
		Rate	43.1	16.6
		Population	2,360,432	2,563,461
2012	1,350	Reported Diagnoses	980	370
		Rate	41.5	14.4
		Population	2,378,589	2,581,035
2013	1,310	Reported Diagnoses	941	369
		Rate	39.6	14.3
		Population	2,393,238	2,597,050
2014	1,256	Reported Diagnoses	892	364
		Rate	37.3	14.0
		Population	2,408,068	2,611,589
2015	1,189	Reported Diagnoses	867	322
		Rate	36.0	12.3
		Population	2,417,553	2,620,599
2016	1,121	Reported Diagnoses	807	314
		Rate	33.4	12.0
		Population	2,427,953	2,631,783
2017	1,050	Reported Diagnoses	757	293
		Rate	31.2	11.1
		Population	2,434,054	2,640,003
2018	1,021	Reported Diagnoses	738	283
		Rate	30.3	10.7
		Population	2,439,706	2,647,094
2019	931	Reported Diagnoses	662	269
		Rate	27.1	10.2


<u>Figure 5 – Trends in Reported HIV Diagnoses, Among Residents at Diagnosis Aged 13+, by Race/Ethnicity, 1985-2019, Reported through June 30, 2020</u>

Number and Percent by Race/Ethnicity of People Diagnosed with HIV, Age 13+ at HIV Diagnosis, by Year of HIV Diagnosis from 1985 through 2019, as Reported through June 30, 2020





Reported HIV Diagnoses Among People Aged 13+ by Race/Ethnicity, 1985-2019, Reported through June 30, 2020




Non-Hispanic Other race not shown

Reported HIV Diagnoses Among People Aged 13+ Race/Ethnicity Year of HIV Non-Hispanic Black, Non-Hispanic White, Non-Hispanic Other Diagnosis No. Hispanic only only Race No. % No. No. No. % <1985 381 11 2.9% 214 56.2% 142 37.3% 14 3.7% 1985 711 20 2.8% 403 56.7% 37.3% 3.2% 265 23 932 558 1986 33 3.5% 59.9% 322 34.5% 19 2.0% 1987 1,237 30 2.4% 762 61.6% 414 33.5% 31 2.5% 2.2% 48 68.0% 453 37 1988 1,679 2.9% 1,141 27.0% 72.9% 1989 2,128 32 1.5% 1,551 498 23.4% 47 2.2% 1990 43 1.9% 1,622 72.8% 511 22.9% 52 2.3% 2,228 1991 2,546 71 2.8% 76.6% 2.2% 1,950 468 18.4% 57 1992 2,394 58 2.4% 1,867 78.0% 405 16.9% 64 2.7% 1993 2,162 64 3.0% 1,704 78.8% 338 15.6% 56 2.6% 1994 1,997 51 2.6% 1,581 79.2% 307 15.4% 58 2.9% 1995 51 15.7% 1,979 2.6% 1,566 79.1% 310 52 2.6% 1996 1,830 48 2.6% 1,484 81.1% 235 12.8% 63 3.4% 1997 50 2.7% 78.4% 268 14.3% 85 1,870 1,467 4.5% 1998 46 2.5% 1,472 246 13.5% 1,827 80.6% 63 3.4% 1999 1,786 65 3.6% 1,406 78.7% 226 12.7% 89 5.0% 1,705 65 3.8% 77.8% 12.4% 102 2000 1,326 212 6.0% 1,442 2001 1,810 61 3.4% 79.7% 213 11.8% 94 5.2% 2002 2,022 67 3.3% 1,607 79.5% 231 11.4% 117 5.8% 2003 1,911 99 5.2% 1,487 77.8% 221 11.6% 104 5.4% 2004 1,949 76 3.9% 1,539 79.0% 232 11.9% 102 5.2% 2005 1,993 96 4.8% 1,554 78.0% 222 11.1% 121 6.1% 2006 2,131 126 5.9% 1,620 76.0% 266 12.5% 119 5.6% 2007 129 6.0% 1,618 75.5% 272 12.7% 125 5.8% 2,144 2008 2,041 147 7.2% 1,498 73.4% 279 13.7% 117 5.7% 2009 6.9% 1,275 75.2% 207 12.2% 97 5.7% 1,696 117 2010 1,743 123 7.1% 72.7% 232 120 1,268 13.3% 6.9% 2011 1,430 132 9.2% 1,024 71.6% 192 13.4% 82 5.7% 108 982 72.7% 103 2012 1,350 8.0% 157 11.6% 7.6% 2013 1,310 95 7.3% 967 73.8% 171 13.1% 77 5.9% 2014 1,256 113 9.0% 923 73.5% 143 11.4% 77 6.1% 2015 103 885 1,189 8.7% 74.4% 134 11.3% 67 5.6% 2016 1,121 92 8.2% 829 74.0% 148 13.2% 52 4.6% 2017 1,050 103 9.8% 747 149 4.9% 71.1% 14.2% 51 2018 1,021 82 8.0% 769 75.3% 126 12.3% 44 4.3% 2019 931 102 11.0% 690 74.1% 102 37 4.0% 11.0% Total 59,490 2,757 4.6% 44,798 75.3% 9,317 15.7% 2,618 4.4%

<u>Figure 5A – Trends in Rates of Reported HIV Diagnoses, Among Residents at Diagnosis</u> <u>Aged 13+, by Race/Ethnicity, 2010-2019, Reported through June 30, 2020</u>


Rate by Race/Ethnicity of People Diagnosed with HIV, Age 13+ at HIV Diagnosis, by Year of HIV Diagnosis from 2010 through 2019, as Reported through June 30, 2020




Year of HIV	Number of			Race/Ethnicity		
Diagnosis	HIV Diagnoses		Hispanic	Non-Hispanic Black	Non-Hispanic White	Non-Hispanic Other Race
		Population	360,847	1,381,323	2,729,639	357,466
2010	1,743	Reported Diagnoses	123	1,268	232	120
		Rate	34.1	91.8	8.5	33.6
		Population	372,819	1,400,333	2,735,168	369,892
2011	1,430	Reported Diagnoses	132	1,024	192	82
		Rate	35.4	73.1	7.0	22.2
		Population	384,916	1,418,066	2,738,538	382,373
2012	1,350	Reported Diagnoses	108	982	157	103
		Rate	28.1	69.2	5.7	26.9
		Population	396,443	1,435,211	2,734,402	393,567
2013	1,310	Reported Diagnoses	95	967	171	77
		Rate	24.0	67.4	6.3	19.6
		Population	409,008	1,452,443	2,723,788	405,049
2014	1,256	Reported Diagnoses	113	923	143	77
		Rate	27.6	63.5	5.3	19.0
		Population	422,464	1,469,819	2,709,950	417,423
2015	1,189	Reported Diagnoses	103	885	134	67
		Rate	24.4	60.2	4.9	16.1
		Population	435,171	1,483,250	2,692,385	427,346
2016	1,121	Reported Diagnoses	92	829	148	52
		Rate	21.1	55.9	5.5	12.2
		Population	449,239	1,497,458	2,674,642	438,396
2017	1,050	Reported Diagnoses	103	747	149	51
		Rate	22.9	49.9	5.6	11.6
		Population	463,686	1,508,218	2,655,979	446,175
2018	1,021	Reported Diagnoses	82	769	126	44
		Rate	17.7	51.0	4.7	9.9
		Population	476,656	1,519,091	2,636,574	454,479
2019	931	Reported Diagnoses	102	690	102	37
		Rate	21.4	45.4	3.9	8.1


<u>Figure 6 – Trends in Reported HIV Diagnoses, Among Residents at Diagnosis Aged 13+, 1985-2019, Reported through June 30, 2020</u>

Number and Percent by Age at HIV Diagnosis of People Diagnosed with HIV, Age 13+ at HIV Diagnosis, by Year of HIV Diagnosis from 1985 through 2019, as Reported through June 30, 2020






Reported HIV Diagnoses Among People Aged 13+ by Age at Diagnosis, 1985-2019, Reported through June 30, 2020



					Reported	HIV Diagno	oses Amoi	ng People	Aged 13+				
Year of HIV						Α	ge at HIV	Diagnosis					
Diagnosis	No.	13-	19	20-	29	30-	39	40-	49	50-	-59	60	+
		No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
<1985	381	33	8.7%	157	41.2%	114	29.9%	54	14.2%	19	5.0%	4	1.0%
1985	711	32	4.5%	283	39.8%	268	37.7%	84	11.8%	24	3.4%	20	2.8%
1986	932	32	3.4%	360	38.6%	350	37.6%	129	13.8%	43	4.6%	18	1.9%
1987	1,237	32	2.6%	427	34.5%	525	42.4%	167	13.5%	60	4.9%	26	2.1%
1988	1,679	22	1.3%	537	32.0%	760	45.3%	258	15.4%	71	4.2%	31	1.8%
1989	2,128	42	2.0%	635	29.8%	976	45.9%	354	16.6%	91	4.3%	30	1.4%
1990	2,228	42	1.9%	650	29.2%	974	43.7%	405	18.2%	109	4.9%	48	2.2%
1991	2,546	44	1.7%	699	27.5%	1,133	44.5%	517	20.3%	109	4.3%	44	1.7%
1992	2,394	45	1.9%	592	24.7%	1,100	45.9%	503	21.0%	112	4.7%	42	1.8%
1993	2,162	34	1.6%	461	21.3%	1,039	48.1%	473	21.9%	120	5.6%	35	1.6%
1994	1,997	24	1.2%	428	21.4%	901	45.1%	489	24.5%	121	6.1%	34	1.7%
1995	1,979	31	1.6%	365	18.4%	872	44.1%	525	26.5%	132	6.7%	54	2.7%
1996	1,830	28	1.5%	305	16.7%	819	44.8%	497	27.2%	133	7.3%	48	2.6%
1997	1,870	28	1.5%	313	16.7%	803	42.9%	526	28.1%	134	7.2%	66	3.5%
1998	1,827	40	2.2%	292	16.0%	750	41.1%	532	29.1%	165	9.0%	48	2.6%
1999	1,786	28	1.6%	257	14.4%	722	40.4%	542	30.3%	182	10.2%	55	3.1%
2000	1,705	47	2.8%	246	14.4%	654	38.4%	513	30.1%	179	10.5%	66	3.9%
2001	1,810	34	1.9%	262	14.5%	648	35.8%	605	33.4%	195	10.8%	66	3.6%
2002	2,022	72	3.6%	331	16.4%	682	33.7%	644	31.8%	219	10.8%	74	3.7%
2003	1,911	56	2.9%	305	16.0%	620	32.4%	593	31.0%	255	13.3%	82	4.3%
2004	1,949	49	2.5%	365	18.7%	576	29.6%	639	32.8%	242	12.4%	78	4.0%
2005	1,993	54	2.7%	380	19.1%	521	26.1%	680	34.1%	289	14.5%	69	3.5%
2006	2,131	65	3.1%	432	20.3%	544	25.5%	675	31.7%	308	14.5%	107	5.0%
2007	2,144	100	4.7%	468	21.8%	505	23.6%	635	29.6%	342	16.0%	94	4.4%
2008	2,041	88	4.3%	496	24.3%	454	22.2%	549	26.9%	308	15.1%	146	7.2%
2009	1,696	89	5.2%	420	24.8%	400	23.6%	459	27.1%	251	14.8%	77	4.5%
2010	1,743	77	4.4%	471	27.0%	371	21.3%	454	26.0%	290	16.6%	80	4.6%
2011	1,430	63	4.4%	442	30.9%	294	20.6%	339	23.7%	219	15.3%	73	5.1%
2012	1,350	80	5.9%	432	32.0%	277	20.5%	300	22.2%	175	13.0%	86	6.4%
2013	1,310	58	4.4%	417	31.8%	299	22.8%	273	20.8%	172	13.1%	91	6.9%
2014	1,256	48	3.8%	420	33.4%	305	24.3%	255	20.3%	164	13.1%	64	5.1%
2015	1,189	49	4.1%	437	36.8%	287	24.1%	205	17.2%	157	13.2%	54	4.5%
2016	1,121	50	4.5%	403	36.0%	262	23.4%	183	16.3%	146	13.0%	77	6.9%
2017	1,050	59	5.6%	373	35.5%	271	25.8%	153	14.6%	124	11.8%	70	6.7%
2018	1,021	58	5.7%	347	34.0%	272	26.6%	152	14.9%	143	14.0%	49	4.8%
2019	931	22	2.4%	325	34.9%	241	25.9%	157	16.9%	110	11.8%	76	8.2%
Total	59,490	1,755	3.0%	14,533	24.4%	20,589	34.6%	14,518	24.4%	5,913	9.9%	2,182	3.7%

<u>Figure 6A – Trends in Rates of Reported HIV Diagnoses, Among Residents at Diagnosis</u> <u>Aged 13+, by Age at Diagnosis, 2010-2019, Reported through June 30, 2020</u>


Rate by Age at HIV Diagnosis of People Diagnosed with HIV, Age 13+ at HIV Diagnosis, by Year of HIV Diagnosis from 2010 through 2019, as Reported through June 30, 2020




Year of HIV	Number of			Age at HI	IV Diagnosis			
Diagnosis	HIV Diagnoses		13-19	20-29	30-39	40-49	50-59	60+
		Population	557,355	790,677	747,128	878,017	823,132	1,032,975
2010	1,743	Reported Diagnoses	77	471	371	454	290	80
		Rate	13.8	59.6	49.7	51.7	35.2	7.7
		Population	554,054	801,210	748,052	868,581	839,856	1,066,033
2011	1,430	Reported Diagnoses	63	442	294	339	219	73
		Rate	11.4	55.2	39.3	39.0	26.1	6.8
		Population	547,765	808,639	756,426	856,555	854,489	1,100,133
2012	1,350	Reported Diagnoses	80	432	277	300	175	86
		Rate	14.6	53.4	36.6	35.0	20.5	7.8
		Population	541,347	814,256	768,133	837,037	865,554	1,133,839
2013	1,310	Reported Diagnoses	58	417	299	273	172	91
		Rate	10.7	51.2	38.9	32.6	19.9	8.0
		Population	536,877	816,094	780,091	814,114	874,562	1,169,458
2014	1,256	Reported Diagnoses	48	420	305	255	164	64
		Rate	8.9	51.5	39.1	31.3	18.8	5.5
		Population	534,707	813,859	792,949	796,574	877,584	1,205,143
2015	1,189	Reported Diagnoses	49	437	287	205	157	54
		Rate	9.2	53.7	36.2	25.7	17.9	4.5
		Population	533,837	807,153	803,766	781,124	873,630	1,240,221
2016	1,121	Reported Diagnoses	50	403	262	183	146	77
		Rate	9.4	49.9	32.6	23.4	16.7	6.2
		Population	533,021	801,406	814,131	771,504	864,841	1,276,124
2017	1,050	Reported Diagnoses	59	373	271	153	124	70
		Rate	11.1	46.5	33.3	19.8	14.3	5.5
		Population	532,906	794,984	823,580	763,740	853,899	1,311,557
2018	1,021	Reported Diagnoses	58	347	272	152	143	49
		Rate	10.9	43.6	33.0	19.9	16.7	3.7
		Population	532,906	794,984	823,580	763,740	853,899	1,311,557
2019	931	Reported Diagnoses	22	325	241	157	110	76
		Rate	4.1	40.9	29.3	20.6	12.9	5.8


<u>Figure 7 – Trends in Reported HIV Diagnoses, Among Residents at Diagnosis Aged 13+, by</u> <u>Estimated Exposure Category, 1985-2019, Reported through June 30, 2020</u>

Number and Percent by Estimated Exposure Category[§] of People Diagnosed with HIV, Age 13+ at HIV Diagnosis, by Year of HIV Diagnosis from 1985 through 2019, as Reported through June 30, 2020





Reported HIV Diagnoses Among People Aged 13+ by Estimated Exposure Category, 1985-2019, Reported through June 30, 2020



Other exposure category not shown

 \S Multiple imputation was used to estimate and adjust for missing exposure category.

Reported HIV Diagnoses Among People Aged 13+ **Estimated Exposure Category§** Year of HIV % with No Diagnosis IDU MSM/IDU No. Reported **MSM HET** Other Exposure Exposure[¥] % No. No. % No. % No. % No. % <1985 381 5.2% 223 58.5% 22.4% 39 10.2% 5.2% 14 3.7% 86 20 1985 711 5.1% 354 49.8% 208 29.2% 78 11.0% 41 5.8% 30 4.2% 1986 932 3.1% 475 51.0% 290 31.1% 78 8.4% 6.7% 26 2.8% 63 36 1987 3.2% 571 107 6.7% 1,237 46.2% 439 35.5% 8.7% 83 2.9% 1988 3.9% 39.4% 41.4% 1,679 661 695 152 9.1% 136 8.1% 35 2.1% 1989 2,128 3.6% 752 953 176 9.6% 35.3% 44.8% 8.3% 205 43 2.0% 4.2% 748 45.2% 177 7.9% 11.9% 1990 2,228 33.6% 1,008 265 30 1.3% 4.8% 1991 2,546 752 29.5% 1,235 48.5% 163 6.4% 362 14.2% 35 1.4% 1992 2,394 6.0% 597 24.9% 1,262 52.7% 161 6.7% 355 14.8% 19 0.8% 1993 4.3% 527 24.4% 51.0% 138 6.4% 17.3% 21 2,162 1,102 375 1.0% 1994 1,997 5.2% 462 23.1% 1,006 50.4% 100 5.0% 409 20.5% 20 1.0% 1995 1,979 6.3% 452 22.8% 949 47.9% 118 6.0% 446 22.5% 14 0.7% 1996 1,830 7.1% 385 21.0% 896 49.0% 105 5.7% 430 23.5% 14 0.8% 400 47.3% 1997 1,870 8.4% 21.4% 884 92 4.9% 488 26.1% 5 0.3% 103 9 1998 1,827 8.6% 348 19.0% 869 47.6% 5.7% 498 27.2% 0.5% 1999 1,786 9.3% 389 21.8% 791 44.3% 83 4.6% 519 29.1% 5 0.3% 2000 1,705 9.9% 360 21.1% 672 39.4% 82 4.8% 588 34.5% 3 0.2% 2001 1,810 10.0% 357 19.7% 752 41.6% 86 4.8% 613 33.9% 2 0.1% 2002 11.2% 443 21.9% 729 71 777 38.4% 2 2,022 36.1% 3.5% 0.1% 2003 1,911 10.5% 458 24.0% 669 35.0% 78 4.1% 701 36.7% 4 0.2% 2004 1,949 13.4% 500 25.7% 30.9% 3.0% 785 40.3% 603 58 4 0.2% 2005 1,993 16.8% 524 26.3% 572 28.7% 72 3.6% 824 41.3% 2 0.1% 2006 2,131 21.4% 584 27.4% 589 27.6% 97 4.5% 857 40.2% 4 0.2% 879 3 2007 2,144 24.3% 695 32.4% 499 23.3% 68 3.2% 41.0% 0.1% 2008 2,041 31.0% 760 37.2% 17.9% 50 2.5% 0.0% 366 865 42.4% 1 731 0 2009 1,696 31.5% 706 41.6% 224 13.2% 35 2.1% 43.1% 0.0% 44.7% 2010 1,743 33.6% 778 207 11.9% 1.9% 722 41.5% 33 1 0.1% 2011 1,430 31.0% 695 48.6% 139 9.7% 33 2.3% 564 39.4% 0 0.0% 2012 1,350 20.6% 682 50.5% 106 7.9% 24 1.7% 538 39.9% 1 0.1% 2013 1,310 16.3% 644 49.2% 96 7.3% 34 534 40.8% 2.6% 1 0.1% 2014 1,256 15.0% 659 52.5% 70 5.5% 16 1.3% 510 40.6% 1 0.1% 2015 1,189 9.7% 629 52.9% 72 6.1% 14 1.2% 473 39.8% 1 0.1% 2016 1,121 14.8% 584 52.1% 66 5.8% 21 1.9% 450 40.1% 1 0.1% 2017 1,050 15.0% 564 53.7% 68 6.5% 16 1.5% 402 38.2% 0.1% 1 0.2% 2018 1,021 15.6% 540 52.9% 61 6.0% 11 1.1% 407 39.9% 2 2019 931 36.9% 477 51.2% 73 7.8% 15 1.6% 39.3% 0.1% 366 1 29.0% 59,490 13.0% 33.2% 32.5% 2,784 4.7% 17,281 391 0.7% Total 19,735 19,306

Note. Discrepancies in the table between total HIV diagnoses and the sums by exposure category are due to less than 1 percent of people missing transmission category after multiple imputation.

[§] Multiple imputation was used to estimate and adjust for missing transmission category.

^{*}Percent with no reported exposure prior to multiple imputation adjustment.


Section III - Continuum of Care Cascades

In their 2011 article in Clinical Infectious Diseases, Gardner et al.¹ utilize a continuum of care "cascade" to illustrate estimates of the number of persons living with HIV who belong to each of the stages of engagement in HIV care.

<u>Figure 8 – Prevalence-Based Estimated 2019 HIV Continuum of Care Cascades, Among</u> Current Maryland Residents Aged 13+, Reported through June 30, 2020

The prevalence-based Maryland Engagement in HIV Care Cascades (below) apply the Centers for Disease Control and Prevention (CDC) CD4 depletion model to Maryland surveillance data, reported through June 30, 2020, to estimate the number of people living with undiagnosed HIV (10.8 percent) and the estimated number of people living with HIV currently residing in Maryland. Maryland surveillance data is then used to describe the number and percentage of persons living with HIV in Maryland on December 31, 2019 and the number and percentage of new HIV diagnoses in 2019 who belong to each of the stages of engagement in HIV care.





Among current Maryland residents living with HIV


Among 2019 HIV diagnoses

Continuum of Care Stage	People Aged 13+	Living with HIV	Continuum of Cara Stage	Reported 2019 HIV Diagnoses			
	Number	Percent	Continuum of Care Stage	Number	Percent		
HIV Infected	35,460	100.0%	New HIV Diagnoses	931	100.0%		
HIV Diagnosed	31,630	89.2%	Linked to HIV Care – 1 Month	809	86.9%		
Retained in HIV Care	25,459	71.8%	Linked to HIV Care – 3 Months	847	91.0%		
Suppressed VL	22,057	62.2%					


¹ Gardner et al, "The Spectrum of Engagement in HIV Care and its Relevance to Test-and-Treat Strategies for Prevention of HIV Infection." Clin Infect Dis. 2011; 52 (6): 793-800.

<u>Figure 8A – Prevalence-Based Estimated 2019 HIV Continuum of Care Cascades, Among Current Maryland Residents Aged</u> 13+, by Sex at Birth, Reported through June 30, 2020


Prevalence-Based Estimated HIV Continuum of Care Among People Aged 13+ by Sex at Birth, 2019




<u>Figure 8B – Prevalence-Based Estimated 2019 HIV Continuum of Care Cascades, Among Current Maryland Residents Aged</u> 13+, by Race/Ethnicity, Reported through June 30, 2020



<u>Figure 8C – Prevalence-Based Estimated 2019 HIV Continuum of Care Cascades, Among Current Maryland Residents</u> Aged 13+, by Age on December 31, 2019, Reported through June 30, 2020






Note. Age groups are consistent with the national standard presentation of the HIV continuum of care by age. A diagnosis-based continuum of care broken down by 10-year age groups is available on the Maryland Department of Health HIV by Age Data Sheet.

<u>Figure 8D – Prevalence-Based Estimated 2019 HIV Continuum of Care Cascades, Among Current Maryland Residents Aged</u> 13+, by Estimated Exposure Category, Reported through June 30, 2020

Prevalence-Based Estimated HIV Continuum of Care Among People Aged 13+ by Estimated Exposure Category, 2019




							Continuun	n of Care Sta	ge						
Demographic -			People Age	d 13+ Livin	g with Diag	nosed HIV			R	Reported 2019 Diagnoses Among People Aged 13+					
2 cm og. apme	HIV In	fected	HIV Diag	HIV Diagnosed Re		Retained in Care Suppre		sed VL	HIV Di	agnosed	Linked to	HIV Care	Linked to	HIV Care	
			2		netanica	care	Зарріс			ивпоэси	- 1 N	lonth	- 3 Months		
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	
Sex at Birth															
Male	23,742	100.0%	20,893	88.0%	16,648	70.1%	14,440	60.8%	662	100.0%	588	88.8%	612	92.4%	
Female	11,734	100.0%	10,737	91.5%	8,811	75.1%	7,617	64.9%	269	100.0%	221	82.2%	235	87.4%	
Age on December 31, 2019															
13-24	1,402	100.0%	862	61.5%	737	52.6%	554	39.5%	173	100.0%	150	86.7%	156	90.2%	
25-34	6,543	100.0%	5,045	77.1%	3,998	61.1%	3,161	48.3%	312	100.0%	268	85.9%	285	91.3%	
35-44	6,884	100.0%	6,030	87.6%	4,675	67.9%	3,987	57.9%	187	100.0%	163	87.2%	169	90.4%	
45-54	8,663	100.0%	8,091	93.4%	6,517	75.2%	5,678	65.5%	135	100.0%	123	91.1%	126	93.3%	
55+	12,149	100.0%	11,602	95.5%	9,532	78.5%	8,677	71.4%	124	100.0%	105	84.7%	111	89.5%	
Race/Ethnicity															
American Indian/Alaska Native	17	100.0%	17	99.9%	9	52.9%	8	47.0%	1	100.0%	1	100.0%	1	100.0%	
Asian	288	100.0%	243	84.5%	182	63.3%	165	57.4%	17	100.0%	17	100.0%	17	100.0%	
Black/African American	26,344	100.0%	23,578	89.5%	19,008	72.2%	16,297	61.9%	690	100.0%	596	86.4%	624	90.4%	
Hispanic/Latino	2,557	100.0%	2,171	84.9%	1,695	66.3%	1,478	57.8%	102	100.0%	89	87.3%	95	93.1%	
Native Hawaiian/Other Pacific Islander	2	100.0%	2	97.4%	2	97.4%	2	97.4%	0	100.0%					
White	4,394	100.0%	3,937	89.6%	3,134	71.3%	2,843	64.7%	102	100.0%	89	87.3%	93	91.2%	
Multiple races	1,886	100.0%	1,682	89.2%	1,429	75.8%	1,264	67.0%	19	100.0%	17	89.5%	17	89.5%	
Exposure Category															
Male-to-male sexual contact															
(MSM)	14,538	100.0%	12,576	86.5%	10,136	69.7%	8,828	60.7%	477	100.0%	423	80.7%	443	88.7%	
Injection drug use (IDU)	5,478	100.0%	5,330	97.3%	4,359	79.6%	3,698	67.5%	73	100.0%	60	71.3%	60	81.3%	
Male	3,166	100.0%	3,071	97.0%	2,441	77.1%	2,090	66.0%	38	100.0%	33	78.2%	33	82.5%	
Female	2,312	100.0%	2,259	97.7%	1,917	82.9%	1,608	69.5%	35	100.0%	27	59.3%	27	79.3%	
MSM/IDU	1,198	100.0%	1,148	95.8%	940	78.4%	800	66.8%	15	100.0%	11	84.2%	13	87.5%	
Heterosexual contact (HET)	13,748	100.0%	12,057	87.7%	9,584	69.7%	8,404	61.1%	366	100.0%	314	86.8%	329	93.2%	
Male	4,632	100.0%	3,863	83.4%	2,939	63.5%	2,577	55.6%	132	100.0%	120	87.4%	121	93.8%	
Female	9,124	100.0%	8,194	89.8%	6,645	72.8%	5,827	63.9%	234	100.0%	194	86.5%	208	92.9%	
Total	35,460	100.0%	31,630	89.2%	25,459	71.8%	22,057	62.2%	931	100.0%	809	86.9%	847	91.0%	

Note. Other exposure category and perinatal transmission not shown. Multiple imputation was used to estimate and adjust for missing transmission category. Exposure category estimates are rounded to whole numbers after percentages are calculated. This may be more noticeable in the precision of estimates calculated for small numbers of people.

<u>Figure 9 – Diagnosis-Based Estimated 2019 HIV Continuum of Care Cascades, Among</u> Current Maryland Residents Aged 13+, Reported through June 30, 2020

The diagnosis-based Maryland Engagement in HIV Care Cascade (below) uses Maryland surveillance data for current Maryland residents to describe the number and percentage of persons living with diagnosed HIV in Maryland on December 31, 2019 and the number and percentage of new HIV diagnoses in 2019 who belong to each of the stages of engagement in HIV care.






Continuum of Care Stage	People Aged 13+	Living with HIV	Continuum of Cara Stage	Reported 2019 HIV Diagnoses			
	Number	Percent	Continuum of Care Stage	Number	Percent		
HIV Diagnosed	31,630	100.0%	New HIV Diagnoses	931	100.0%		
Retained in HIV Care	25,459	80.5%	Linked to HIV Care – 1 Month	809	86.9%		
Suppressed VL	22,068	69.8%	Linked to HIV Care – 3 Months	847	91.0%		

Section IV - By Jurisdiction

Maryland Reported HIV Diagnoses Among People Aged 13+, Rate by Jurisdiction, 2019



• The rate of new HIV diagnoses among people aged 13+ in Maryland 18.6 is per 100,000.

Jurisdictions with the highest rate of new HIV diagnoses among people aged 13+ in Maryland include:

Baltimore City: 39.7 per 100,000
 Prince George's County: 38.0 per 100,000
 Charles County: 25.7 per 100,000
 Wicomico County: 22.9 per 100,000

• The rate of new AIDS diagnoses among people aged 13+ in Maryland is **10.6 per 100,000**.

Jurisdictions with the highest rate of new AIDS diagnoses among people aged 13+ in Maryland include:

Baltimore City: 29.9 per 100,000
 Kent County: 17.3 per 100,000
 Prince George's County: 17.2 per 100,000
 Wicomico County: 13.7 per 100,000

<u>Table 1 – Reported HIV Diagnoses and Reported AIDS Diagnoses, Among People Aged 13+</u> <u>During 2019, by Jurisdiction of Residence at Diagnosis, Reported through June 30, 2020</u>

Jurisdiction of Residence at	Population Age	Reported HIV I	Diagnoses Among 13+	People Aged	Reported AIDS Diagnoses Among People Aged 13+				
Diagnosis	13+	No.	% of Total	Rate	No.	% of Total	Rate		
Allegany	61,764	3	0.3%	4.9	3	0.6%	4.9		
Anne Arundel	485,680	48	5.2%	9.9	15	3.1%	3.1		
Baltimore City	504,269	200	21.5%	39.7	151	31.5%	29.9		
Baltimore	698,716	117	12.6%	16.7	49	10.2%	7.0		
Calvert	77,905	3	0.3%	3.9	3	0.6%	3.9		
Caroline	27,787	3	0.3%	10.8	0	0.0%	0.0		
Carroll	143,123	5	0.5%	3.5	6	1.3%	4.2		
Cecil	86,690	7	0.8%	8.1	0	0.0%	0.0		
Charles	136,026	35	3.8%	25.7	11	2.3%	8.1		
Dorchester	27,165	2	0.2%	7.4	0	0.0%	0.0		
Frederick	217,096	13	1.4%	6.0	0	0.0%	0.0		
Garrett	25,324	0	0.0%	0.0	0	0.0%	0.0		
Harford	215,720	10	1.1%	4.6	6	1.3%	2.8		
Howard	270,649	23	2.5%	8.5	8	1.7%	3.0		
Kent	17,295	3	0.3%	17.3	3	0.6%	17.3		
Montgomery	877,406	135	14.5%	15.4	72	15.0%	8.2		
Prince George's	761,035	289	31.0%	38.0	131	27.3%	17.2		
Queen Anne's	42,995	3	0.3%	7.0	0	0.0%	0.0		
Saint Mary's	94,126	5	0.5%	5.3	0	0.0%	0.0		
Somerset	22,450	0	0.0%	0.0	0	0.0%	0.0		
Talbot	32,407	0	0.0%	0.0	0	0.0%	0.0		
Washington	127,813	5	0.5%	3.9	5	1.0%	3.9		
Wicomico	87,334	20	2.1%	22.9	12	2.5%	13.7		
Worcester	46,024	2	0.2%	4.3	1	0.2%	2.2		
Corrections		0	0.0%		4	0.8%			
Total	5,086,800	931	100.0%	18.6	480	100.0%	10.6		

Table 2 – Reported HIV Diagnoses, Among People Aged 13+ During 2019, Linked to Care, Late Diagnosis, First CD4 Test Result, and Mean Years from HIV Diagnosis by Jurisdiction of Residence at Diagnosis, Reported through June 30, 2020

	Reported HIV Diagnoses Among People Aged 13+										
Jurisdiction of — Residence at HIV		Linked to	Care	% Late HIV –	First CD4 Test Result						
Diagnosis	No.	% 1 month	% 3 months	Diagnoses	% with Test	Median Count					
Allegany	3	***	***	***	***	***					
Anne Arundel	48	81.3%	85.4%	18.8%	87.5%	371					
Baltimore City	200	86.0%	88.5%	24.0%	85.5%	369					
Baltimore	117	90.6%	94.9%	18.8%	93.2%	423					
Calvert	3	***	***	***	***	***					
Caroline	3	***	***	***	***	***					
Carroll	5	***	100.0%	***	100.0%	282					
Cecil	7	71.4%	71.4%	0.0%	71.4%	525					
Charles	35	80.0%	88.6%	20.0%	94.3%	334					
Dorchester	2	***	***	***	***	***					
Frederick	13	53.8%	61.5%	0.0%	69.2%	326					
Garrett	0										
Harford	10	80.0%	90.0%	***	90.0%	330					
Howard	23	91.3%	95.7%	21.7%	95.7%	304					
Kent	3	***	***	***	***	***					
Montgomery	135	90.4%	94.8%	33.3%	92.6%	287					
Prince George's	289	87.9%	91.7%	21.8%	92.0%	395					
Queen Anne's	3	***	***	***	***	***					
Saint Mary's	5	***	***	0.0%	100.0%	342					
Somerset	0										
Talbot	0										
Washington	5	***	100.0%	***	100.0%	616					
Wicomico	20	90.0%	90.0%	30.0%	90.0%	445					
Worcester	2	***	***	***	***	***					
Corrections	0										
Total	931	86.9%	91.0%	22.9%	90.4%	377					

^{***} Data withheld due to low population counts and/or small numbers

<u>Table 3 – People Aged 13+ Living with Diagnosed HIV, Alive on December 31, 2019, by</u>
<u>Jurisdiction of Residence at Diagnosis and Current Residence, Reported through</u>
<u>June 30, 2020</u>

		Total People Aged 13+ Living with Diagnosed HIV										
Jurisdiction of Residence	Population		Residence a	at Diagnosis			%					
	Age 13+	No.	% of Total	Rate	Ratio (1 in X)	No.	% of Total	Rate	Ratio (1 in X)	% Change		
Allegany	61,764	82	0.2%	132.8	753	112	0.4%	181.3	551	36.6%		
Anne Arundel	485,680	1,294	3.9%	266.4	375	1,398	4.4%	287.8	347	8.0%		
Baltimore City	504,269	11,964	36.2%	2,372.5	42	10,600	33.5%	2,102.1	48	-11.4%		
Baltimore	698,716	3,210	9.7%	459.4	218	3,769	11.9%	539.4	185	17.4%		
Calvert	77,905	114	0.3%	146.3	683	127	0.4%	163.0	613	11.4%		
Caroline	27,787	68	0.2%	244.7	409	66	0.2%	237.5	421	-2.9%		
Carroll	143,123	153	0.5%	106.9	935	162	0.5%	113.2	883	5.9%		
Cecil	86,690	122	0.4%	140.7	711	142	0.4%	163.8	610	16.4%		
Charles	136,026	494	1.5%	363.2	275	616	1.9%	452.9	221	24.7%		
Dorchester	27,165	131	0.4%	482.2	207	148	0.5%	544.8	184	13.0%		
Frederick	217,096	368	1.1%	169.5	590	454	1.4%	209.1	478	23.4%		
Garrett	25,324	10	0.0%	39.5	2,532	17	0.1%	67.1	1,490	70.0%		
Harford	215,720	463	1.4%	214.6	466	516	1.6%	239.2	418	11.4%		
Howard	270,649	629	1.9%	232.4	430	695	2.2%	256.8	389	10.5%		
Kent	17,295	38	0.1%	219.7	455	46	0.1%	266.0	376	21.1%		
Montgomery	877,406	4,091	12.4%	466.3	214	3,544	11.2%	403.9	248	-13.4%		
Prince George's	761,035	7,779	23.5%	1,022.2	98	7,926	25.1%	1,041.5	96	1.9%		
Queen Anne's	42,995	52	0.2%	120.9	827	57	0.2%	132.6	754	9.6%		
Saint Mary's	94,126	141	0.4%	149.8	668	160	0.5%	170.0	588	13.5%		
Somerset	22,450	60	0.2%	267.3	374	104	0.3%	463.2	216	73.3%		
Talbot	32,407	65	0.2%	200.6	499	75	0.2%	231.4	432	15.4%		
Washington	127,813	301	0.9%	235.5	425	349	1.1%	273.1	366	15.9%		
Wicomico	87,334	250	0.8%	286.3	349	264	0.8%	302.3	331	5.6%		
Worcester	46,024	76	0.2%	165.1	606	67	0.2%	145.6	687	-11.8%		
Corrections		1,108	3.4%			216	0.7%					
Total	5,086,800	33,063	100.0%	650.0	154	31,630	100.0%	621.8	161	-4.3%		

Table 4 – CD4 Test Results During 2019 for People Aged 13+ Living with Diagnosed HIV, Alive on December 31, 2019, by Jurisdiction of Current Residence, Reported through June 30, 2020

	Total People Aged 13+ Living with Diagnosed HIV										
Jurisdiction of Current				Recer	nt CD4 Test Re	sult					
Residence	No.	No. with Test	% with Test	Median Count	<200	200-349	350-499	500+			
Allegany	112	104	92.9%	618	9.6%	6.7%	19.2%	64.4%			
Anne Arundel	1,398	1,060	75.8%	627	6.0%	12.5%	16.8%	64.6%			
Baltimore City	10,600	8,027	75.7%	592	10.0%	12.7%	16.4%	61.0%			
Baltimore	3,769	2,869	76.1%	647	7.4%	10.7%	14.6%	67.2%			
Calvert	127	100	78.7%	686	8.0%	7.0%	17.0%	68.0%			
Caroline	66	60	90.9%	694	1.7%	11.7%	5.0%	81.7%			
Carroll	162	117	72.2%	688	6.8%	11.1%	12.8%	69.2%			
Cecil	142	90	63.4%	606	10.0%	6.7%	18.9%	64.4%			
Charles	616	487	79.1%	640	7.0%	12.5%	14.2%	66.3%			
Dorchester	148	119	80.4%	647	4.2%	16.0%	14.3%	65.5%			
Frederick	454	356	78.4%	628	3.7%	9.0%	16.9%	70.5%			
Garrett	17	14	82.4%	670	0.0%	0.0%	14.3%	85.7%			
Harford	516	381	73.8%	582	7.9%	13.6%	15.7%	62.7%			
Howard	695	530	76.3%	643	7.7%	9.1%	15.7%	67.5%			
Kent	46	41	89.1%	683	4.9%	7.3%	4.9%	82.9%			
Montgomery	3,544	2,569	72.5%	605	7.2%	11.3%	16.7%	64.8%			
Prince George's	7,926	6,108	77.1%	617	7.3%	11.3%	16.3%	65.1%			
Queen Anne's	57	39	68.4%	729	10.3%	5.1%	12.8%	71.8%			
Saint Mary's	160	123	76.9%	591	4.9%	21.1%	15.4%	58.5%			
Somerset	104	93	89.4%	619	6.5%	10.8%	12.9%	69.9%			
Talbot	75	65	86.7%	524	12.3%	15.4%	18.5%	53.8%			
Washington	349	274	78.5%	644	7.3%	8.8%	16.4%	67.5%			
Wicomico	264	217	82.2%	562	12.0%	13.8%	17.1%	57.1%			
Worcester	67	56	83.6%	628	3.6%	10.7%	17.9%	67.9%			
Corrections	216	181	83.8%	647	3.3%	9.9%	19.9%	66.9%			
Total	31,630	24,080	76.1%	614	8.1%	11.7%	16.1%	64.1%			

<u>Table 5 – Viral Load Test Results During 2019 for People Aged 13+ Living with Diagnosed</u>
<u>HIV, Alive on December 31, 2019, by Jurisdiction of Current Residence, Reported through</u>
<u>June 30, 2020</u>

<u> </u>	Total People Aged 13+ Living with Diagnosed HIV									
Jurisdiction of Current Residence	_		Recent Viral Lo	ad Test Result						
	No.	No. with Test	% with Test	% Suppressed	Median Unsuppressed					
Allegany	112	103	92.0%	91.3%	3,610					
Anne Arundel	1,398	1,092	78.1%	89.5%	14,900					
Baltimore City	10,600	8,525	80.4%	86.3%	13,357					
Baltimore	3,769	3,030	80.4%	90.2%	6,934					
Calvert	127	101	79.5%	93.1%	5,560					
Caroline	66	57	86.4%	96.5%	1,560					
Carroll	162	122	75.3%	91.0%	7,160					
Cecil	142	94	66.2%	85.1%	3,995					
Charles	616	494	80.2%	90.1%	6,820					
Dorchester	148	119	80.4%	93.3%	6,895					
Frederick	454	358	78.9%	92.7%	1,657					
Garrett	17	13	76.5%	100.0%						
Harford	516	390	75.6%	90.0%	7,870					
Howard	695	545	78.4%	89.9%	7,907					
Kent	46	42	91.3%	81.0%	6,350					
Montgomery	3,544	2,586	73.0%	91.0%	10,868					
Prince George's	7,926	6,183	78.0%	88.9%	7,500					
Queen Anne's	57	43	75.4%	90.7%	8,220					
Saint Mary's	160	127	79.4%	88.2%	11,900					
Somerset	104	94	90.4%	89.4%	3,785					
Talbot	75	63	84.0%	90.5%	6,395					
Washington	349	276	79.1%	87.7%	8,535					
Wicomico	264	218	82.6%	85.3%	18,640					
Worcester	67	56	83.6%	94.6%	15,000					
Corrections	216	177	81.9%	86.4%	544					
Total	31,630	24,908	78.7%	88.6%	10,000					

Section V - By Age

<u>Table 6 – Reported HIV Diagnoses During 2019, Linked to Care, Late Diagnosis, and First</u> CD4 Test Result by Age at HIV Diagnosis, Reported through June 30, 2020

		Reported HIV Diagnoses									
Age at HIV Diagnosis	Population	Na		Rate -	Linked	to Care	Late HIV Diagnoses	First CD4 Test Result			
Diagnosis	nosis No. % of Total Rate % %	%	%	% with	Median						
					1 month	3 months	70	Test	Count		
<5 (Pediatric)	361,937	0	0.0%	0.0							
5-12 (Pediatric)	596,943	0	0.0%	0.0							
13-19	530,841	22	2.4%	4.1	86.4%	86.4%	***	86.4%	367		
20-29	783,259	325	34.9%	41.5	85.2%	90.5%	12.3%	89.8%	425		
30-39	827,633	241	25.9%	29.1	88.4%	92.1%	23.7%	91.7%	373		
40-49	757,459	157	16.9%	20.7	87.3%	91.1%	31.2%	89.2%	299		
50-59	841,381	110	11.8%	13.1	88.2%	90.0%	32.7%	90.0%	307		
60+	1,346,227	76	8.2%	5.6	86.8%	92.1%	***	93.4%	308		
Total	6,045,680	931	100.0%	15.4	86.9%	91.0%	22.9%	90.4%	377		

^{***} Data withheld due to low population counts and/or small numbers

<u>Table 7 – People Living with Diagnosed HIV, Alive on December 31, 2019, by Age on December 31, 2019, Reported through June 30, 2020</u>

Age on	Total People Living with Diagnosed HIV								
December 31, 2019	Population	No. % of Total		Rate	Ratio (1 in X)				
<5 (Pediatric)	361,937	6	0.0%	1.7	60,323				
5-12 (Pediatric)	596,943	44	0.1%	7.4	13,567				
13-19	530,841	158	0.5%	29.8	3,360				
20-29	783,259	2,747	8.7%	350.7	285				
30-39	827,633	6,002	18.9%	725.2	138				
40-49	757,459	6,542	20.7%	863.7	116				
50-59	841,381	9,448	29.8%	1,122.9	89				
60+	1,346,227	6,733	21.3%	500.1	200				
Total	6,045,680	31,680	100.0%	524.0	191				

Table 8 – CD4 Test Results During 2019 for People Aged 13+ Living with Diagnosed HIV, Alive on December 31, 2019, by Age on December 31, 2019, Reported through June 30, 2020

	Total People Aged 13+ Living with Diagnosed HIV										
Age on	Recent CD4 Test Results										
December 31, 2019	No.	No. with Test	% with Test	Median Count	<200	200-349	350-499	500+			
13-19	158	135	85.4%	658	2.2%	7.4%	16.3%	74.1%			
20-29	2,747	2,162	78.7%	636	7.8%	8.8%	15.9%	67.5%			
30-39	6,002	4,442	74.0%	626	8.3%	10.4%	15.9%	65.3%			
40-49	6,542	4,890	74.7%	618	8.5%	12.1%	15.2%	64.2%			
50-59	9,448	7,290	77.2%	612	8.4%	12.2%	15.8%	63.6%			
60+	6,733	5,161	76.7%	594	7.2%	13.2%	17.6%	62.1%			
Total	31,630	24,080	76.1%	614	8.1%	11.7%	16.1%	64.1%			

<u>Table 9 – Viral Load Test Results During 2019 for People Living with Diagnosed HIV, Alive on December 31, 2019, by Age on December 31, 2019, Reported through June 30, 2020</u>

	Total People Aged 13+ Living with Diagnosed HIV									
Age on			Recent Viral Load Test Result							
December 31, 2019	No.	No. with Test	% with Test	% Suppressed	Median Unsuppressed					
13-19	158	141	89.2%	82.3%	4,080					
20-29	2,747	2,212	80.5%	77.7%	13,900					
30-39	6,002	4,587	76.4%	83.7%	11,212					
40-49	6,542	5,026	76.8%	89.1%	11,282					
50-59	9,448	7,574	80.2%	90.4%	8,260					
60+	6,733	5,368	79.7%	94.2%	4,930					
Total	31,630	24,908	78.7%	88.6%	10,000					

Section VI – By Demographics

<u>Table 10 – Reported HIV Diagnoses, Among People Aged 13+ During 2019, Linked to Care, Late Diagnosis, and First CD4 Test Result by Sex at Birth, Gender, Race/Ethnicity, and Country of Birth, Reported through June 30, 2020</u>

				Reported	d HIV Diagno	ses Among P	eople Aged 13-	+	
Demographic Characteristics	Population Age 13+		% of		Linked	to Care	Late HIV Diagnoses	First CD4 1	est Result
	Age 13+	No.	Total	Rate	% 1 month	% 3 months	%	% with Test	Median Count
Sex at Birth									
Male	2,439,706	662	71.1%	27.1	88.8%	92.4%	20.5%	90.5%	367
Female	2,647,094	269	28.9%	10.2	82.2%	87.4%	28.6%	90.3%	384
Gender									
Male	2,439,706	648	69.6%	26.6	88.9%	92.6%	21.0%	90.4%	357
Female	2,647,094	269	28.9%	10.2	82.2%	87.4%	28.6%	90.3%	384
Transfeminine		14	1.5%	0.0	85.7%	85.7%	0.0%	92.9%	717
Transmasculine		0	0.0%	0.0					
Another Gender Identity		0	0.0%	0.0					
Race/Ethnicity									
Hispanic	476,656	102	11.0%	21.4	87.3%	93.1%	30.4%	92.2%	312
Non-Hispanic	4,610,144	829	89.0%	18.0	86.9%	90.7%	22.0%	90.2%	381
American Indian/Alaska Native, only	12,702	1	0.1%	7.9	***	***	***	***	***
Asian, only	341,287	17	1.8%	5.0	100.0%	100.0%	***	94.1%	390
Black, only	1,519,091	690	74.1%	45.4	86.4%	90.4%	21.7%	90.3%	382
Native Hawaiian/Other Pacific Islander, only	2,519	0	0.0%	0.0					
White, only	2,636,574	102	11.0%	3.9	87.3%	91.2%	23.5%	89.2%	357
Multiracial/Other	97,971	19	2.0%	19.4	***	***	***	***	***
Total	5,086,800	931	100.0%	18.3	86.9%	91.0%	22.9%	90.4%	377
Country of Birth									
United States	4,172,343	683	73.4%	16.4	87.6%	91.7%	19.5%	90.6%	384
Foreign-Born	863,070	154	16.5%	17.8	86.4%	90.9%	35.7%	91.6%	310
Africa		92	9.9%		85.9%	91.3%	31.5%	92.4%	390
Asia		7	0.8%		100.0%	100.0%	***	100.0%	317
Caribbean		13	1.4%		69.2%	76.9%	61.5%	76.9%	71
Central America		28	3.0%		89.3%	92.9%	42.9%	92.9%	212
Other		14	1.5%		92.9%	92.9%	***	92.9%	300
Unknown		94	10.1%		83.0%	86.2%	26.6%	87.2%	321
Total	5,035,413	931	100.0%	18.5	86.9%	91.0%	22.9%	90.4%	377

Note. Population data by sex at birth, age, and race/ethnicity are from the 2019 U.S. Census population estimates. Population data by gender are unavailable. Population data by country of birth are from the 2018 American Community Survey.

^{***} Data withheld due to low population counts and/or small numbers

<u>Table 11 – Reported HIV Diagnoses, Among People Aged 13+ During 2019, by Sex at Birth and Race/Ethnicity, Reported through June 30, 2020</u>

			Reported H	IV Diagnoses	Among People A	ged 13+					
	Sex at Birth										
Dago /Ethnisity		Male				Fema	ale				
Race/Ethnicity	Population 13+	No.	% of Total	Rate	Population 13+	No.	% of Total	Rate			
Hispanic	247,093	80	12.1%	32.4	229,563	22	8.2%	9.6			
Non-Hispanic	2,192,613	582	87.9%	26.5	2,417,531	247	91.8%	26.5			
American Indian/Alaska Native, only	6,118	1	0.2%	16.3	6,584	0	0.0%	0.0			
Asian, only	159,506	14	2.1%	8.8	181,781	3	1.1%	1.7			
Black, only	696,819	476	71.9%	68.3	822,272	214	79.6%	26.0			
Native Hawaiian/Other Pacific Islander, only	1,176	0	0.0%	0.0	1,343	0	0.0%	0.0			
White, only	1,283,046	78	11.8%	6.1	1,353,528	24	8.9%	1.8			
Multiracial/Other	45,948	13	2.0%	28.3	52,023	6	2.2%	11.5			
Total	2,439,706	662	100.0%	27.1	2,647,094	269	100.0%	27.1			

Note. Population data by sex at birth, age, and race/ethnicity are from the 2019 U.S. Census population estimates.

<u>Table 12 – People Aged 13+ Living with Diagnosed HIV Alive on December 31, 2019, by Sex at Birth, Gender, Race/Ethnicity, and Country of Birth, Reported through June 30, 2020</u>

	Population —	Total P	eople Aged 13+ Livi	ng with Diagnose	d HIV
Demographic Characteristics	Age 13+	No.	% of Total	Rate	Ratio (1 in X)
Sex at Birth					
Male	2,439,706	20,893	66.1%	856.4	117
Female	2,647,094	10,737	33.9%	405.6	247
Gender					
Male	2,439,706	20,579	65.1%	843.5	119
Female	2,647,094	10,717	33.9%	404.9	247
Transfeminine		313	1.0%		
Transmasculine		20	0.1%		
Another Gender Identity		1	0.0%		
Race/Ethnicity					
Hispanic	476,656	2,171	6.9%	455.5	220
Non-Hispanic	4,610,144	29,459	93.1%	639.0	156
American Indian/Alaska Native, only	12,702	17	0.1%	133.8	747
Asian, only	341,287	243	0.8%	71.2	1,404
Black, only	1,519,091	23,578	74.5%	1,552.1	64
Native Hawaiian/Other Pacific Islander, only	2,519	2	0.0%	79.4	1,260
White, only	2,636,574	3,937	12.4%	149.3	670
Multiracial/Other	97,971	1,682	5.3%	1,716.8	58
Total	5,086,800	31,630	100.0%	621.8	161
Country of Birth					
United States	4,172,343	26,979	85.3%	646.6	155
Foreign-Born	863,070	3,680	11.6%	426.4	235
Africa		2,315	7.3%		
Asia		208	0.7%		
Caribbean		282	0.9%		
Central America		581	1.8%		
Other		294	0.9%		
Unknown		971	3.1%		
Total	5,035,413	31,630	100.0%	628.2	159

Note. Population data by sex at birth, age, and race/ethnicity are from the 2019 U.S. Census population estimates. Population data by gender are unavailable. Population data by country of birth are from the 2018 American Community Survey.

<u>Table 13 – People Aged 13+ Living with Diagnosed HIV Alive on December 31, 2019, by Sex at Birth and Race/Ethnicity, Reported through June 30, 2020</u>

		Total People Aged 13+ Living with Diagnosed HIV									
	Sex at Birth										
Dogo/Ethnicity		Male				Fema	ile				
Race/Ethnicity	Population 13+	No.	% of Total	Rate	Population 13+	No.	% of Total	Rate			
Hispanic	247,093	1,668	8.0%	675.0	229,563	503	4.7%	219.1			
Non-Hispanic	2,192,613	19,225	92.0%	876.8	2,417,531	10,234	95.3%	423.3			
American Indian/Alaska Native, only	6,118	11	0.1%	179.8	6,584	6	0.1%	91.1			
Asian, only	159,506	198	0.9%	124.1	181,781	45	0.4%	24.8			
Black, only	696,819	14,685	70.3%	2,107.4	822,272	8,893	82.8%	1,081.5			
Native Hawaiian/Other Pacific Islander, only	1,176	2	0.0%	170.0	1,343	0	0.0%	0.0			
White, only	1,283,046	3,172	15.2%	247.2	1,353,528	765	7.1%	56.5			
Multiracial/Other	45,948	1,157	5.5%	2,518.1	52,023	525	4.9%	1,009.2			
Total	2,439,706	20,893	100.0%	856.4	2,647,094	10,737	100.0%	405.6			

Note. Population data by sex at birth, age, and race/ethnicity are from the 2019 U.S. Census population estimates.

Table 14 – CD4 Test Results During 2019 for People Aged 13+ Living with Diagnosed HIV, Alive on December 31, 2019, by Sex at Birth, Gender, Race/Ethnicity and Country of Birth, Reported through June 30, 2020

	Total People Aged 13+ Living with Diagnosed HIV									
Demographic Characteristics				Recen	t CD4 Test R	esult				
Demographic characteristics	No.	No. with Test	% with Test	Median Count	<200	200-349	350-499	500+		
Sex at Birth										
Male	20,893	15,790	75.6%	595	8.1%	12.3%	17.2%	62.5%		
Female	10,737	8,290	77.2%	654	8.1%	10.7%	14.1%	67.2%		
Gender										
Male	20,579	15,546	75.5%	594	8.1%	12.3%	17.2%	62.4%		
Female	10,717	8,273	77.2%	653	8.1%	10.7%	14.1%	67.1%		
Transfeminine	313	243	77.6%	640	7.8%	9.5%	15.2%	67.5%		
Transmasculine	20	***	***	***	***	***	***	***		
Another Gender Identity	1	***	***	***	***	***	***	***		
Race/Ethnicity										
Hispanic	2,171	1,627	74.9%	598	8.2%	12.0%	16.8%	62.9%		
Non-Hispanic	29,459	22,453	76.2%	615	8.1%	11.7%	16.0%	64.2%		
American Indian/Alaska Native, only	17	***	***	***	***	***	***	***		
Asian, only	243	177	72.8%	585	5.1%	11.3%	19.8%	63.8%		
Black, only	23,578	18,003	76.4%	610	8.3%	11.9%	16.3%	63.4%		
Native Hawaiian/Other Pacific Islander, only	2	***	***	***	***	***	***	***		
White, only	3,937	2,915	74.0%	661	6.7%	9.9%	14.4%	69.0%		
Multiracial/Other	1,682	1,348	80.1%	603	7.9%	12.8%	15.4%	63.9%		
Country of Birth										
United States	26,979	20,710	76.8%	619	8.2%	11.6%	15.8%	64.3%		
Foreign-Born	3,680	2,683	72.9%	585	7.1%	12.8%	18.2%	61.9%		
Africa	2,315	1,678	72.5%	574	7.0%	13.0%	19.7%	60.4%		
Asia	208	156	75.0%	597	5.1%	10.3%	20.5%	64.1%		
Caribbean	282	224	79.4%	662	12.1%	8.5%	12.1%	67.4%		
Central America	581	423	72.8%	576	7.1%	13.9%	17.5%	61.5%		
Other	294	202	68.7%	621	4.0%	15.3%	12.9%	67.8%		
Unknown	971	687	70.8%	629	7.4%	10.3%	16.6%	65.6%		
Total	31,630	24,080	76.1%	614	8.1%	11.7%	16.1%	64.1%		

^{***} Data withheld due to low population counts and/or small numbers

Table 15 – Viral Load Test Results During 2019 for People Aged 13+ Living with Diagnosed HIV, Alive on December 31, 2019, by Sex at Birth, Gender, Race/Ethnicity and Country of Birth, Reported through June 30, 2020

	Total People Aged 13+ Living with Diagnosed HIV									
Demographic Characteristics			Recent Viral Lo	ad Test Result						
Demograpme characteristics	No.	No. with Test	% with Test	% Suppressed	Median Unsuppressed					
Sex at Birth										
Male	20,893	16,296	78.0%	88.6%	10,884					
Female	10,737	8,612	80.2%	88.4%	8,390					
Gender										
Male	20,579	16,043	78.0%	88.8%	10,733					
Female	10,717	8,594	80.2%	88.4%	8,390					
Transfeminine	313	252	80.5%	79.0%	13,600					
Transmasculine	20	***	***	***	**					
Another Gender Identity	1	***	***	***	**:					
Race/Ethnicity										
Hispanic	2,171	1,660	76.5%	89.0%	13,70					
Non-Hispanic	29,459	23,248	78.9%	88.5%	9,92					
American Indian/Alaska Native, only	17	***	***	***	**					
Asian, only	243	178	73.3%	92.7%	23,60					
Black, only	23,578	18,604	78.9%	87.6%	10,00					
Native Hawaiian/Other Pacific Islander, only	2	***	***	***	**					
White, only	3,937	3,051	77.5%	93.2%	9,52					
Multiracial/Other	1,682	1,404	83.5%	90.0%	8,70					
Country of Birth										
United States	26,979	21,476	79.6%	88.2%	10,500					
Foreign-Born	3,680	2,726	74.1%	91.9%	4,69					
Africa	2,315	1,702	73.5%	91.4%	3,89					
Asia	208	156	75.0%	94.9%	13,04					
Caribbean	282	228	80.9%	92.5%	38,41					
Central America	581	434	74.7%	91.7%	4,26					
Other	294	206	70.1%	93.7%	7,87					
Unknown	971	706	72.7%	87.5%	9,74					
Total	31,630	24,908	78.7%	88.6%	10,000					

^{***} Data withheld due to low population counts and/or small numbers

<u>Table 16 – Males Living with Diagnosed HIV by Age on December 31, 2019 and Race/Ethnicity, Alive on December 31, 2019, and Reported through June 30, 2020</u>

Total Males Living with Diagnosed HIV									
A			Race/E	thnicity		Total Males			
Age on December 31, 2019		Hispanic	Non-Hispanic Black, only	Non-Hispanic White, only	Non-Hispanic Other Race	by Age Group			
	Population	85,637	147,147	199,490	56,885	489,159			
442 (D - - -	Number	3	15	2	0	20			
<13 (Pediatric)	Rate	3.5	10.2	1.0	0.0	4.1			
	Ratio (1 in X)	28,546	9,810	99,745		24,458			
	Population	37,035	83,460	119,788	29,989	270,272			
12.10	Number	5	76	6	2	89			
13-19	Rate	13.5	91.1	5.0	6.7	32.9			
	Ratio (1 in X)	7,407	1,098	19,965	14,995	3,037			
	Population	47,156	128,825	182,522	37,660	396,163			
20.20	Number	204	1,704	151	104	2,163			
20-29	Rate	432.6	1,322.7	82.7	276.2	546.0			
	Ratio (1 in X)	231	76	1,209	362	183			
	Population	56,662	118,264	192,602	38,865	406,393			
20.20	Number	450	3,232	426	288	4,396			
30-39	Rate	794.2	2,732.9	221.2	741.0	1,081.7			
	Ratio (1 in X)	126	37	452	135	92			
	Population	50,346	107,840	173,117	35,730	367,033			
40.40	Number	447	2,475	579	278	3,779			
40-49	Rate	887.9	2,295.1	334.5	778.1	1,029.6			
	Ratio (1 in X)	113	44	299	129	97			
	Population	31,826	116,491	224,304	29,738	402,359			
50.50	Number	379	4,119	1,138	418	6,054			
50-59	Rate	1,190.9	3,535.9	507.3	1,405.6	1,504.6			
	Ratio (1 in X)	84	28	197	71	66			
	Population	24,068	141,939	390,713	40,766	597,486			
60	Number	183	3,079	872	278	4,412			
60+	Rate	760.3	2,169.2	223.2	681.9	738.4			
	Ratio (1 in X)	132	46	448	147	135			
	Population	332,730	843,966	1,482,536	269,633	2,928,865			
Total Males by	Number	1,671	14,700	3,174	1,368	20,913			
Race/Ethnicity	Rate	502.2	1,741.8	214.1	507.4	714.0			
	Ratio (1 in X)	199	57	467	197	140			

Table 17 – Females Living with Diagnosed HIV by Age on December 31, 2019 and Race/Ethnicity, Alive on December 31, 2019, and Reported through June 30, 2020

		Total Fema	ales Living with Diag	nosed HIV		
Agoon			Race/E	thnicity		Total Females
Age on December 31, 2019		Hispanic	Non-Hispanic Black, only	Non-Hispanic White, only	Non-Hispanic Other Race	by Age Group
	Population	81,529	144,029	189,717	54,447	469,721
412 (Dadiatuia)	Number	1	25	2	2	30
<13 (Pediatric)	Rate	1.2	17.4	1.1	3.7	6.4
	Ratio (1 in X)	81,529	5,761	94,859	27,223	15,657
	Population	35,077	82,009	114,041	29,441	260,569
12.10	Number	3	61	2	3	69
13-19	Rate	8.6	74.4	1.8	10.2	26.5
	Ratio (1 in X)	11,692	1,344	57,021	9,814	3,776
	Population	43,380	131,112	174,005	38,599	387,096
20.20	Number	40	485	33	26	584
20-29	Rate	92.2	369.9	19.0	67.4	150.9
	Ratio (1 in X)	1,085	270	5,273	1,485	663
	Population	49,013	135,212	192,136	44,879	421,240
20.20	Number	94	1,277	138	97	1,606
30-39	Rate	191.8	944.4	71.8	216.1	381.3
	Ratio (1 in X)	521	106	1,392	463	262
	Population	43,153	127,505	177,472	42,296	390,426
40-49	Number	161	2,249	191	162	2,763
40-49	Rate	373.1	1,763.9	107.6	383.0	707.7
	Ratio (1 in X)	268	57	929	261	141
	Population	29,601	140,052	233,773	35,596	439,022
50.50	Number	134	2,853	253	154	3,394
50-59	Rate	452.7	2,037.1	108.2	432.6	773.1
	Ratio (1 in X)	221	49	924	231	129
	Population	29,339	206,382	462,101	50,919	748,741
CO :	Number	71	1,968	148	134	2,321
60+	Rate	242.0	953.6	32.0	263.2	310.0
	Ratio (1 in X)	413	105	3,122	380	323
	Population	311,092	966,301	1,543,245	296,177	3,116,815
Total Females by	Number	504	8,918	767	578	10,767
Race/Ethnicity	Rate	162.0	922.9	49.7	195.2	345.4
	Ratio (1 in X)	617	108	2,012	512	289
-						

Section VII - By HIV Exposure Category

<u>Table 18 – Reported HIV Diagnoses, Among People Aged 13+ During 2019, Linked to Care, Late Diagnosis, and First CD4 Test Result, by Estimated or Reported Exposure Category and Sex at Birth, Reported through June 30, 2020</u>

	Reported HIV Diagnoses Among People Aged 13+								
Estimated or Reported Exposure Category	NI-	0/ -f.T-t-1	Linked	to Care	Late HIV Diagnoses	First CD4 Test Result			
Exposure Category	No.	% of Total -	% 1 month	% 3 months	%	% with Test	Median Count		
Male-to-male Sexual Contact (MSM)	477	51.2%	88.7%	92.9%	16.6%	91.5%	383		
Injection Drug Use (IDU)	73	7.8%	82.2%	83.3%	30.2%	85.2%	317		
Male-to-male Sexual Contact and Injection Drug Use (MSM/IDU)	15	1.6%	***	***	***	***	***		
Heterosexual Contact (HET)	366	39.3%	85.9%	89.9%	29.6%	90.1%	358		
Perinatal Transmission	1	0.1%	***	***	***	***	***		
Other Exposure	0	0.0%							
No Estimated or Reported Exposure	0	0.0%							
Total	931	100.0%	86.9%	91.0%	22.9%	90.4%	377		
Exposure Category Male-to-male Sexual Contact (MSM) Injection Drug Use (IDU)	477	72.1% 5.7%	88.7% 86.9%	92.9%	16.6% 33.6%	91.5%	383		
							383		
Male-to-male Sexual Contact and Injection	36	5.7%	80.9%	88.8%	33.0%	80.7%	320		
Drug Use (MSM/IDU)	15	2.2%	***	***	***	***	***		
Heterosexual Contact (HET)	132	19.9%	91.1%	91.8%	30.9%	88.0%	298		
Perinatal Transmission	1	0.2%	***	***	***	***	***		
Other Exposure	0	0.0%							
No Estimated or Reported Exposure	0	0.0%							
Total	662	100.0%	88.8%	92.4%	20.5%	90.5%	367		
Female Estimated or Reported Exposure Category									
Injection Drug Use (IDU)	35	13.0%	77.1%	77.4%	26.6%	83.7%	295		
Heterosexual Contact (HET)	234	87.0%	82.9%	88.8%	28.9%	91.3%	396		
Perinatal Transmission	0	0.0%							
Other Exposure	0	0.0%							
No Estimated or Reported Exposure	0	0.0%							
Total	269	100.0%	82.2%	87.4%	28.6%	90.3%	384		

Note. Data for 2019 are preliminary because they are based on a 6-month reporting delay and should not be used to assess trends. Data have been statistically adjusted to account for missing transmission category, therefore, values may not sum to column total. Exposure category estimates are rounded to whole numbers after percentages are calculated. This may be more noticeable in the precision of estimates calculated for small numbers of people. Median count is weighted based on statistical adjustment. For exposure category by gender, please see the Maryland HIV Epidemiological Profile Gender Diverse People Supplement.

^{***} Data withheld due to low population counts and/or small numbers

<u>Table 19 – People Aged 13+ Living with Diagnosed HIV, Alive on December 31, 2019, by Estimated or Reported Exposure Category and Sex at Birth, Reported through June 30, 2020</u>

Estimated on Devented European Cotons	Total People Aged 13+ Living with Diagnosed HIV			
Estimated or Reported Exposure Category	No.	% of Total		
Male-to-male Sexual Contact (MSM)	12,576	39.8%		
Injection Drug Use (IDU)	5,330	16.9%		
Male-to-male Sexual Contact and Injection Drug Use (MSM/IDU)	1,148	3.6%		
Heterosexual Contact (HET)	12,057	38.1%		
Perinatal Transmission	398	1.3%		
Other Exposure	45	0.1%		
No Estimated or Reported Exposure	77	0.2%		
Total	31,630	100.0%		
Male Estimated or Reported Exposure Category				
Male-to-male Sexual Contact (MSM)	12,576	60.2%		
Injection Drug Use (IDU)	3,071	14.7%		
Male-to-male Sexual Contact and Injection Drug Use (MSM/IDU)	1,148	5.5%		
Heterosexual Contact (HET)	3,863	18.5%		
Perinatal Transmission	170	0.8%		
Other Exposure	24	0.1%		
No Estimated or Reported Exposure	42	0.2%		
Total	20,893	100.0%		
Female Estimated or Reported Exposure Category				
Injection Drug Use (IDU)	2,259	21.0%		
Heterosexual Contact (HET)	8,194	76.3%		
Perinatal Transmission	228	2.1%		
Other Exposure	21	0.2%		
No Estimated or Reported Exposure	35	0.3%		
Total	10,737	100.0%		

Note. Data have been statistically adjusted to account for missing transmission category, therefore, values may not sum to column total. For exposure category by gender, please see the Maryland HIV Epidemiological Profile Gender Diverse People Supplement.

<u>Table 20 – People Aged 13+ Living with Diagnosed HIV by Estimated or Reported Exposure</u>
<u>Category, Race/Ethnicity, and Sex at Birth, Alive on December 31, 2019, and Reported</u>
<u>through June 30, 2020</u>

		Total Ped	ple Aged 13+ L	iving with Diagnose	ed HIV	
Estimated or Reported			Race/E	thnicity		
Exposure Category	His	panic	Non-Hispan	ic, Black, only	Non-Hispanic, White, only	
	No.	% of Total	No.	% of Total	No.	% of Total
Male-to-male Sexual Contact (MSM)	1,118	51.5%	8,111	34.4%	2,435	61.8%
Injection Drug Use (IDU)	145	6.7%	4,307	18.3%	566	14.4%
Male-to-male Sexual Contact and Injection Drug Use (MSM/IDU)	65	3.0%	809	3.4%	188	4.8%
Heterosexual Contact (HET)	819	37.7%	9,922	42.1%	708	18.0%
Perinatal Transmission	16	0.7%	346	1.5%	18	0.5%
Other Exposure	2	0.1%	21	0.1%	17	0.4%
No Estimated or Reported Exposure	6	0.3%	62	0.3%	6	0.2%
Total	2,171	100.0%	23,578	100.0%	3,938	100.0%
Male Estimated or Reported Exposure Category						
Male-to-male Sexual Contact (MSM)	1,118	67.0%	8,111	55.2%	2,435	76.8%
Injection Drug Use (IDU)	86	5.2%	2,563	17.5%	271	8.6%
Male-to-male Sexual Contact and Injection Drug Use (MSM/IDU)	65	3.9%	809	5.5%	188	5.9%
Heterosexual Contact (HET)	387	23.2%	3,018	20.5%	253	8.0%
Perinatal Transmission	7	0.4%	146	1.0%	8	0.3%
Other Exposure	2	0.1%	6	0.0%	14	0.4%
No Estimated or Reported Exposure	3	0.2%	33	0.2%	3	0.1%
Total	1,668	100.0%	14,686	100.0%	3,172	100.0%
Female Estimated or Reported Exposure Category						
Injection Drug Use (IDU)	59	11.6%	1,745	19.6%	295	38.5%
Heterosexual Contact (HET)	433	86.0%	6,904	77.6%	454	59.4%
Perinatal Transmission	9	1.8%	200	2.2%	10	1.3%
Other Exposure	0	0.0%	15	0.2%	3	0.4%
No Estimated or Reported Exposure	3	0.6%	29	0.3%	3	0.4%
Total	504	100.0%	8,893	100.0%	765	100.0%

Note. Data have been statistically adjusted to account for missing transmission category, therefore, values may not sum to column total. For exposure category by gender, please see the Maryland HIV Epidemiological Profile Gender Diverse People Supplement.

<u>Table 21 – CD4 Test Results During 2019 for People Aged 13+ Living with Diagnosed HIV, Alive on December 31, 2019, by Estimated or Reported Exposure Category, Reported through June 30, 2020</u>

			Total Peop	le Aged 13+ w	ith Living Dia	agnosed HIV			
Estimated or Reported	Recent CD4 Test Results								
Exposure Category	No.	No. with Test	% with Test	Median Count	<200	200-349	350-499	500+	
Male-to-male Sexual Contact (MSM)	12,576	9,623	76.5%	632	6.3%	10.6%	15.9%	67.2%	
Injection Drug Use (IDU)	5,330	4,047	75.9%	560	12.5%	14.2%	16.5%	56.8%	
Male-to-male Sexual Contact and Injection Drug Use (MSM/IDU)	1,148	890	77.6%	552	9.3%	13.0%	20.4%	57.3%	
Heterosexual Contact (HET)	12,057	9,102	75.5%	624	7.5%	11.9%	15.6%	65.0%	
Perinatal Transmission	398	325	81.7%	583	18.2%	7.7%	15.1%	59.1%	
Other Exposure	45	31	68.9%	474	19.4%	9.7%	22.6%	48.4%	
No Estimated or Reported Exposure	77	62	80.5%	564	9.7%	4.8%	24.2%	61.3%	
Total	31,630	24,080	76.1%	614	8.1%	11.7%	16.1%	64.1%	

Note. Data have been statistically adjusted to account for missing transmission category, therefore, values may not sum to column total. Median count is weighted based on statistical adjustment.

Table 22 – Viral Load Test Results During 2019 for People Aged 13+ Living with Diagnosed HIV, Alive on December 31, 2019, by Estimated or Reported Exposure Category, Reported through June 30, 2020


		Total People	Aged 13+ with Living	g Diagnosed HIV				
Estimated or Reported		Recent Viral Load Test Results						
Exposure Category	No.	No. with Test	% with Test	% Suppressed	Median Unsuppressed			
Male-to-male Sexual Contact (MSM)	12,576	9,942	79.1%	88.8%	11,900			
Injection Drug Use (IDU)	5,330	4,245	79.6%	87.1%	11,500			
Male-to-male Sexual Contact and Injection Drug Use (MSM/IDU)	1,148	920	80.1%	87.0%	11,500			
Heterosexual Contact (HET)	12,057	9,367	77.7%	89.7%	7,920			
Perinatal Transmission	398	337	84.7%	73.6%	10,000			
Other Exposure	45	35	77.8%	88.6%	13,539			
No Estimated or Reported Exposure	77	63	81.8%	76.2%	1,680			
Total	31,630	24,908	78.7%	88.6%	10,000			


Note. Data have been statistically adjusted to account for missing transmission category, therefore, values may not sum to column total. Median unsuppressed is weighted based on statistical adjustment.

Section VIII - Disease Progression and Mortality

<u>Figure 10 – Trends in Disease Progression by Year of HIV Diagnosis, 1985-2018, Reported through June 30, 2020</u>

Number of People Aged 13+ with Diagnosed HIV, Age 13+ at HIV Diagnosis, (Reported HIV Diagnoses Among People Aged 13+), Number of People Aged 13+ with Diagnosed HIV with an AIDS Diagnosis by December 31, 2019 (Reported AIDS Diagnoses), Percent by Year of HIV Diagnosis of Reported HIV Diagnoses with a Reported AIDS Diagnosis Among People Aged 13+ by December 31, 2019, the Average Years from HIV Diagnosis to AIDS Diagnosis for People with an AIDS Diagnosis, and the Percent by Year of HIV Diagnosis of Reported HIV Diagnoses with a Reported AIDS Diagnosis Among People Aged 13+ at 1 Year, 2 Years, 5 Years, and 10 Years after HIV Diagnosis, by Year of HIV Diagnosis from 1985 through 2018, Reported through June 30, 2020







_		Reported HIV Diagnoses Among People Aged 13+							
Year of HIV		R	eported AIDS Diag		Percent	with AIDS by	Years After HIV	Diagnosis	
Diagnosis	No.	No.	% with AIDS	Mean Years from HIV to AIDS	1 Year	2 Years	5 Years	10 Years	
<1985	381	365	95.8%	7.3	37.3%	37.3%	40.4%	65.4%	
1985	711	662	93.1%	6.3	31.1%	32.6%	40.6%	75.2%	
1986	932	884	94.8%	5.3	36.2%	37.2%	49.0%	82.1%	
1987	1,237	1,156	93.5%	4.5	38.0%	40.5%	57.4%	83.6%	
1988	1,679	1,566	93.3%	3.9	40.4%	44.4%	65.6%	86.0%	
1989	2,128	1,950	91.6%	3.5	41.0%	45.9%	69.0%	84.9%	
1990	2,228	2,039	91.5%	3.2	43.9%	51.0%	72.5%	85.4%	
1991	2,546	2,290	89.9%	2.8	46.5%	55.6%	74.0%	84.1%	
1992	2,394	2,171	90.7%	2.8	49.5%	58.4%	74.9%	83.9%	
1993	2,162	1,958	90.6%	2.2	55.7%	63.9%	77.3%	86.0%	
1994	1,997	1,781	89.2%	2.3	57.2%	64.4%	75.6%	83.9%	
1995	1,979	1,737	87.8%	2.4	58.1%	63.7%	72.6%	82.4%	
1996	1,830	1,543	84.3%	2.1	59.0%	63.1%	70.9%	79.7%	
1997	1,870	1,527	81.7%	2.2	55.6%	60.2%	68.6%	76.6%	
1998	1,827	1,482	81.1%	2.1	56.2%	60.0%	68.0%	75.9%	
1999	1,786	1,398	78.3%	1.9	56.3%	60.2%	67.5%	74.0%	
2000	1,705	1,354	79.4%	2.0	54.9%	58.8%	67.9%	75.4%	
2001	1,810	1,420	78.5%	1.9	56.1%	59.7%	68.5%	74.3%	
2002	2,022	1,517	75.0%	1.9	54.2%	57.9%	64.8%	70.9%	
2003	1,911	1,397	73.1%	1.7	53.7%	57.0%	63.0%	69.8%	
2004	1,949	1,379	70.8%	1.9	50.4%	53.8%	59.7%	67.2%	
2005	1,993	1,271	63.8%	1.8	45.8%	48.8%	54.7%	61.2%	
2006	2,131	1,208	56.7%	2.0	38.3%	40.7%	47.8%	54.0%	
2007	2,144	1,093	51.0%	2.2	32.2%	34.9%	41.8%	48.7%	
2008	2,041	988	48.4%	1.9	31.3%	34.5%	41.9%	47.2%	
2009	1,696	778	45.9%	1.9	29.6%	32.3%	39.1%	45.5%	
2010	1,743	739	42.4%	1.7	27.8%	30.6%	36.7%		
2011	1,430	618	43.2%	1.4	30.6%	34.1%	38.3%		
2012	1,350	534	39.6%	1.1	29.6%	32.2%	37.0%		
2013	1,310	502	38.3%	0.8	31.3%	32.8%	36.8%		
2014	1,256	421	33.5%	0.8	25.8%	28.3%	32.8%		
2015	1,189	376	31.6%	0.6	25.9%	28.2%			
2016	1,121	329	29.3%	0.5	24.8%	26.9%			
2017	1,050	308	29.3%	0.3	27.2%	28.8%			
2018	1,021	270	26.4%	0.2	25.3%				
Total	58,559	41,011	70.0%	2.5	43.6%	47.7%	57.3%	65.9%	

<u>Figure 11 – Trends in Survival Times by Year of AIDS Diagnosis, 1985-2018, Reported through June 30, 2020</u>

Number of People Aged 13+ with Diagnosed HIV, Age 13+ at HIV Diagnosis, with an AIDS Diagnosis by December 31, 2018 (Reported AIDS Diagnoses), Number of Reported AIDS Diagnosis Among People Aged 13+ Reported to Have Died of Any Cause by December 31, 2019 (Reported AIDS Deaths), Percent by Year of AIDS Diagnosis of Reported AIDS Diagnoses with a Reported AIDS Death (Percent Dead) Among People Aged 13+ by December 31, 2019, the Average Years from AIDS Diagnosis to Death for People that have Died, and the Percent by Year of AIDS Diagnosis of Reported AIDS Diagnoses Among People Aged 13+ Not Reported to Have Died (Percent Alive) at 1 Year, 2 Years, 5 Years, and 10 Years after AIDS Diagnosis, by Year of AIDS Diagnosis from 1985 through 2018, Reported through June 30, 2020






_	Reported AIDS Diagnoses Among People Aged 13+								
Year of AIDS	_		Reported AIDS D		Perce	nt Alive by Years A	After AIDS Diagno	sis	
Diagnosis	No.	No.	% Deceased	Mean Years from AIDS to Death	1 Year	2 Years	5 Years	10 Years	
<1985	145	140	96.6%	1.7	42.8%	20.7%	10.3%	7.6%	
1985	212	204	96.2%	1.7	42.5%	19.8%	11.3%	6.1%	
1986	314	303	96.5%	2.2	51.3%	33.1%	13.1%	8.3%	
1987	496	486	98.0%	2.2	57.1%	31.7%	11.1%	5.8%	
1988	693	671	96.8%	2.3	60.6%	38.1%	14.0%	6.5%	
1989	931	890	95.6%	2.9	64.0%	43.6%	17.8%	9.8%	
1990	1,173	1,110	94.6%	3.1	66.7%	47.9%	21.2%	12.1%	
1991	1,469	1,364	92.9%	2.9	68.4%	48.2%	18.9%	12.8%	
1992	1,953	1,802	92.3%	3.3	71.8%	50.5%	22.3%	16.3%	
1993	2,257	1,982	87.8%	4.0	77.2%	56.5%	33.1%	22.4%	
1994	2,177	1,819	83.6%	4.2	78.0%	59.0%	39.8%	26.6%	
1995	2,160	1,672	77.4%	5.4	78.8%	67.6%	51.9%	37.6%	
1996	1,936	1,404	72.5%	5.9	82.2%	75.6%	60.1%	43.3%	
1997	1,683	1,131	67.2%	6.2	85.4%	78.3%	63.7%	48.7%	
1998	1,534	998	65.1%	6.1	87.0%	80.8%	66.2%	49.4%	
1999	1,531	939	61.3%	6.0	87.5%	81.1%	66.8%	52.1%	
2000	1,418	792	55.9%	5.9	88.2%	83.6%	69.7%	56.4%	
2001	1,535	877	57.1%	5.5	86.6%	79.6%	66.6%	55.0%	
2002	1,574	822	52.2%	5.7	88.9%	83.9%	71.9%	59.9%	
2003	1,519	727	47.9%	4.8	85.8%	80.4%	70.6%	61.1%	
2004	1,484	630	42.5%	4.9	88.8%	84.4%	75.1%	65.6%	
2005	1,425	572	40.1%	4.9	89.3%	85.4%	76.6%	67.9%	
2006	1,295	453	35.0%	4.2	89.6%	85.8%	77.6%	68.7%	
2007	1,075	368	34.2%	4.2	89.4%	86.4%	79.4%	69.2%	
2008	1,047	334	31.9%	3.6	89.3%	85.4%	78.5%	70.1%	
2009	911	268	29.4%	3.4	90.3%	87.3%	79.9%	71.0%	
2010	1,011	269	26.6%	3.4	91.7%	88.9%	82.8%		
2011	817	187	22.9%	3.2	93.9%	90.9%	83.0%		
2012	877	181	20.6%	2.6	91.4%	89.6%	83.7%		
2013	855	163	19.1%	2.2	92.5%	89.1%	83.4%		
2014	639	91	14.2%	1.5	92.5%	89.7%	86.5%		
2015	628	95	15.1%	1.4	92.0%	89.8%			
2016	593	95	16.0%	1.0	90.6%	87.0%			
2017	585	72	12.3%	0.9	93.2%	89.4%			
2018*	509	41	8.1%	0.2	92.1%				
Total	40,461	23,952	59.2%	4.3	82.6%	73.0%	58.6%	48.8%	

^{*2018} AIDS deaths are preliminary

Section IX – Pediatrics

<u>Figure 12 – Trends in Reported HIV Diagnoses Among People Aged <13, 1985-2019, Reported through June 30, 2020</u>

Number of Pediatric HIV Diagnoses, Age <13 at HIV Diagnosis, (Pediatric Reported HIV Diagnoses) by Year of Birth, by Year of HIV Diagnosis, by Year of AIDS Diagnosis, and by Year of Death Due to Any Cause, and Number of Children with Diagnosed HIV, Age <13 at HIV Diagnosis, and Not Reported to have Died as of December 31st of Each Year (Total Living Pediatric Diagnoses) from 1985 through 2019, as Reported by Name through June 30, 2020



		Pediatric Reporte	d HIV Diagnoses		Total Living Pediatric
Year	No. by Year of Birth	No. by Year of HIV Diagnosis	No. by Year of AIDS Diagnosis	No. by Year of Death	Diagnoses Alive on December 31st of Each Year
<1985	55	5	3	0	
1985	23	13	8	2	17
1986	29	15	6	4	28
1987	32	22	14	5	45
1988	45	28	19	6	67
1989	40	37	25	8	96
1990	65	57	24	9	144
1991	58	59	35	12	191
1992	43	65	39	12	242
1993	53	56	35	16	281
1994	39	41	29	16	306
1995	27	29	24	20	313
1996	27	37	27	19	328
1997	21	18	12	7	339
1998	21	20	12	6	352
1999	11	24	13	5	369
2000	16	16	5	1	384
2001	19	14	11	5	393
2002	16	14	8	4	403
2003	6	6	3	6	402
2004	8	11	6	5	409
2005	7	8	5	1	416
2006	7	14	7	4	427
2007	10	44	10	5	466
2008	9	15	8	3	480
2009	4	10	7	2	489
2010	2	3	8	6	488
2011	2	3	8	5	488
2012	1	3	3	4	490
2013	2	3	11	3	491
2014	3	6	4	1	497
2015	0	0	8	3	495
2016	3	4	6	4	496
2017	2	3	3	3	498
2018	0	3	9	3	498
2019	0	0	6	2	499
Total	706	706	461	217	

<u>Table 23 – People Aged <13 Living with Diagnosed HIV, Alive on December 31, 2019, by</u> <u>Jurisdiction of Current Residence, Reported through June 30, 2020</u>

Lucialistics of Commont Bookless	Total People Aged <13 Living with	n Diagnosed HIV
Jurisdiction of Current Residence	No.	% of Total
Allegany	2	4.0%
Anne Arundel	1	2.0%
Baltimore City	11	22.0%
Baltimore	5	10.0%
Calvert	0	0.0%
Caroline	0	0.0%
Carroll	1	2.0%
Cecil	3	6.0%
Charles	4	8.0%
Dorchester	0	0.0%
Frederick	0	0.0%
Garrett	0	0.0%
Harford	2	4.0%
Howard	1	2.0%
Kent	0	0.0%
Montgomery	7	14.0%
Prince George's	11	22.0%
Queen Anne's	0	0.0%
Saint Mary's	0	0.0%
Somerset	0	0.0%
Talbot	0	0.0%
Washington	2	4.0%
Wicomico	0	0.0%
Worcester	0	0.0%
Corrections	0	0.0%
Total	50	100.0%

<u>Table 24 – People Aged <13 Living with Diagnosed HIV, Alive on December 31, 2019, by Sex at Birth and Race/Ethnicity, Reported through June 30, 2020</u>

Daws a man his Chausatauistis	Total People Aged <13 Living with Diagnosed HIV				
Demographic Characteristic ———	No.	% of Total			
Sex at Birth					
Male	20	40.0%			
Female	30	60.0%			
Race/Ethnicity					
Hispanic	4	8.0%			
Non-Hispanic	46	92.0%			
American Indian/Alaska Native, only	0	0.0%			
Asian, only	1	2.0%			
Black, only	40	80.0%			
Native Hawaiian/Other Pacific Islander, only	0	0.0%			
White, only	4	8.0%			
Multiracial/Other	1	2.0%			
Total	50	100.0%			

Table 25 – Perinatal HIV Transmissions by Year of Birth, 2010-2019

Year of Birth	Live Births	Women of Childbearing Age Living with Diagnosed HIV	Reported Perinatal HIV Exposures	Confirmed P	erinatal HIV Transmi	ssion
	No.	No.	No.	No.	%	Rate
2010	74,999	6,652	177	4	2.3%	5.4
2011	73,783	6,337	174	2	1.1%	2.7
2012	73,052	6,633	183	1	0.5%	1.4
2013	72,751	6,703	163	1	0.6%	1.4
2014	71,806	6,277	202	3	1.5%	4.1
2015	73,588	6,397	152	0	0.0%	0.0
2016	73,544	6,016	185	3	1.6%	4.1
2017	73,073	5,661	169	1	0.6%	1.4
2018	71,589	5,169	162	0	0.0%	0.0
2019		5,008	159	0	0.0%	0.0
Total	658,185	60,853	1,726	15	0.9%	2.3

Live births from Maryland Vital Statistics Annual Report 2019.

Women of childbearing age (ages 13-49) with diagnosed HIV and perinatal HIV exposures from Maryland Enhanced HIV/AIDS Reporting System, reported through June 30, 2020.

Perinatal HIV exposures and transmissions are based on residence at birth.

Glossary of Terms

Age at AIDS Diagnosis: Age group at time of initial AIDS diagnosis.

Age at HIV Diagnosis: Age group at time of initial HIV diagnosis.

Age on [date]: Age group of people living with diagnosed HIV on the specified date.

Another Gender Identity: All other gender identities or gender expressions.

CD4 Result Distribution (<200, 200-349, 350-499, 500+): Percent of people living with diagnosed HIV with a CD4 test distributed by their CD4 count results (cells per microliter).

CD4 With Test: Number and percent of total people aged 13+ living with diagnosed HIV with a recent CD4 test result.

Confirmed Perinatal HIV Transmission: Reported infants born to a woman living with diagnosed HIV, with a confirmed HIV diagnosis and a resident of Maryland at time of birth.

Corrections: Residence in a state or federal prison. Does not include local jails and detention centers.

Current Residence: Jurisdiction of residence from the most recent report since January 1, 2009.

First CD4 Test Result Median Count: Median CD4 count (cells per microliter) of the first CD4 test result reported within 12 months following initial HIV diagnosis.

First CD4 Test Result Percent with Test: Percent of reported HIV diagnoses among people aged 13+ with the first CD4 test result reported within 12 months following the initial HIV diagnosis.

Gender Expression: People's outward presentation of their gender which may or may not conform to socially defined behaviors and characteristics typically associated with being either masculine or feminine.

Gender Identity: Refers to one's internal understanding of one's own gender, or the gender with which a person identifies.

Heterosexual Contact (HET): Men or women who report a history of sexual contact with a person of the opposite sex who is known to be HIV positive or at increased risk for HIV transmission. Risk groups include bisexual men, IDUs and recipients of blood, blood components, transplants of organ or tissue, and artificial insemination.

Injection Drug Use (IDU): Men or women who report receiving an injection, either self-administered or given by another person, of a drug that was not prescribed by a physician.

Jurisdiction of Current Residence: Jurisdiction of residence from the most recent report since January 1, 2009.

Jurisdiction of Residence: Jurisdiction of residence at diagnosis or current residence.

Jurisdiction of Residence at AIDS Diagnosis: Jurisdiction of residence at time of initial AIDS diagnosis.

Jurisdiction of Residence at Diagnosis: Jurisdiction of residence at the later time of initial HIV diagnosis or time of initial AIDS diagnosis.

Jurisdiction of Residence at HIV Diagnosis: Jurisdiction of residence at time of initial HIV diagnosis.

Late HIV Diagnosis: Percent of adult/adolescent reported HIV diagnoses with an initial AIDS diagnosis less than or equal to 3 months after their initial HIV diagnosis.

Linked to Care: Percent of adult/adolescent reported HIV diagnoses with a reported CD4 or viral load test performed less than or equal to 1 month or 3 months after their initial HIV diagnosis.

Live Births: Infants born alive in the general population.

People Aged 13+ Living with Diagnosed HIV: Reported HIV diagnoses, age 13 years or older as of December 31st of the specified year and not reported to have died as of December 31st of the specified year.

People Living with Diagnosed HIV: Reported HIV diagnoses not reported to have died as of December 31st of the specified year.

Mean Years from HIV Diagnosis: Mean or average number of years from initial HIV diagnosis to initial AIDS diagnosis for people living with a reported AIDS diagnosis.

Mean Years from AIDS diagnosis to AIDS Death: Mean or average number of years from initial AIDS diagnosis to death for adult/adolescent reported AIDS diagnoses with a reported death.

Median Count: Median CD4 count (cells per microliter), among total people aged 13+ living with diagnosed HIV, of the most recent CD4 test result measured in the specified year.

Median Unsuppressed: Median unsuppressed viral load (copies per milliliter) among people aged 13+ living with diagnosed HIV of the most recent viral load test result measured in the specified year of 200 copies per milliliter or greater.

Male-to-male Sexual Contact (MSM): Men who report a history of sexual contact with other men (homosexual contact) including men who report sexual contact with both men and women (bisexual contact).

Male-to-male Sexual Contact/Injection Drug Use (MSM/IDU): Men who report a history of sexual contact with other men who also have engaged in injection drug use.

Other Exposure: Includes men or women who received clotting factor for hemophilia/coagulation disorder, or received a transfusion of blood/blood components, or received a transplant of tissue/organs or artificial insemination, or worked in a health care or clinical laboratory setting.

Percent Change: The percent change in number of total people aged 13+ living with diagnosed HIV from residence at diagnosis to current residence.

Percent Alive by Years after AIDS Diagnosis: Percent of adult/adolescent reported AIDS diagnoses not reported to have died at 1 year, 2 years, 5 years and 10 years after initial AIDS diagnosis, by year of AIDS diagnosis.

Percent Late HIV Diagnosis: Percent of adult/adolescent reported AIDS diagnoses with an initial HIV diagnosis less than or equal to 3 months prior to their initial AIDS diagnosis.

Percent Suppressed: Percent of total people aged 13+ living with diagnosed HIV with a recent viral load test result measured in the specified year of less than 200 copies per milliliter.

Percent with AIDS by Years after HIV Diagnosis: Percent of adult/adolescent reported HIV diagnoses in the specified year with an initial AIDS diagnosis at 1 year, 2 years, 5 years and 10 years after HIV diagnosis.

Perinatal Exposure: Any baby born to a woman living with diagnosed HIV.

Perinatal HIV Transmission Percent: Percent of confirmed perinatal HIV transmissions among reported perinatal HIV exposures by year of birth.

Perinatal HIV Transmission Rate: Number of confirmed perinatal HIV transmissions divided by the number of live births multiplied by 100,000 by year of birth.

Population: Population estimate for July 1, 2019.

Population Age 13+: Population age 13 years or older, estimate for July 1, 2019.

Rate: Number of people living with diagnosed HIV divided by the population and multiplied by 100,000.

Ratio (1 in X): Number of people for every 1 person living with diagnosed HIV in the population, or 1 person living with diagnosed HIV in every X number of people.

Recent CD4 Test Result: The most recent CD4 test result measured in the specified year.

Recent Viral Load Test Result: The most recent viral load test result measured in the specified year.

Reported AIDS Deaths: Reported HIV diagnoses with an AIDS diagnosis, reported to have died of any cause during the specified year.

Reported AIDS Diagnoses: Reported HIV diagnoses with an initial AIDS diagnosis during the specified year.

Reported AIDS Diagnoses Among People Aged 13+: Reported HIV diagnoses, age 13 years or older at HIV diagnosis, with an initial AIDS diagnosis during the specified year.

Reported HIV Diagnoses: Reported HIV diagnoses with an initial HIV diagnosis during the specified year.

Reported HIV Diagnoses Among People Aged 13+: Reported HIV diagnoses, age 13 years or older at HIV diagnosis, with an initial HIV diagnosis during the specified year.

Reported Perinatal HIV Exposures: Reported infants born to a woman living with diagnosed HIV during the specified year, with a residence of Maryland at time of birth.

Residence at Diagnosis: Jurisdiction of residence at later time of initial HIV diagnosis or initial AIDS diagnosis.

Retained in HIV Care: HIV diagnosed with a reported CD4 or viral load test result performed or reported antiretroviral use in the specified year.

Suppressed Viral Load: HIV diagnosed with the most recent viral load measured in the specified year of less than 200 copies per milliliter.

Total People Aged 13+ Living with Diagnosed HIV: Reported HIV diagnoses, age 13 years or older as of December 31st of the specified year not reported to have died as of December 31st of the specified year.

Total People Aged <13 Living with Diagnosed HIV: Reported HIV diagnosis, age less than 13 years at HIV diagnosis and not reported to have died as of December 31st of the specified year.

Total People Living with Diagnosed HIV: Reported HIV diagnoses not reported to have died as of December 31st of the specified year.

Transfeminine: A person whose sex assigned at birth is male but who identify as feminine.

Transmasculine: A person whose sex assigned at birth is female but who identify as masculine.

Viral Load With Test: Number and percent of total people aged 13+ living with diagnosed HIV with a recent viral load test result.

Women of Childbearing Age Living with Diagnosed HIV: Reported HIV diagnoses among women of childbearing age (13-49 years old).